

Anomalous soil gas CO₂ concentrations and relation to seismic activity at Rabaul caldera, Papua New Guinea

Niveles de concentración anómalos de CO₂ en los suelos y su relación con la actividad sísmica de la caldera de Rabaúl, Papua Nueva Guinea

N. M. Pérez (*,**), H. Wakita (*), D. Lolok (***), H. Patia (***), B. Talai (***), and C. O. McKee (***)

(*) Laboratory for Earthquake Chemistry THE UNIVERSITY OF TOKYO, Bunkyo-ku, Tokyo 113, Japan
(**) TERRANOSTRA Research Institute, P.O. Box 225, 38400 Puerto de la Cruz, Tenerife, Canary Islands, Spain
(***) Rabaul Volcano Observatory GEOLOGICAL SURVEY OF P.N.G. P.O. Box 386, Rabaul, Papua New Guinea

Resumen

Medidas de CO₂ en los suelos de la caldera de Rabaul, Papua Nueva Guinea, reflejan niveles de concentración altos, hasta cerca de un 20%, en relación con un margen activo de la caldera definido por sismicidad. Estos niveles anómalos del CO₂ en los suelos fueron detectados lejos de los aparatos volcánicos activos. La caracterización isotópica de carbono refleja un origen principalmente biogénico para el CO₂, pero aquellas muestras que presentan un mayor contenido de CO₂ muestran una firma isotópica relativamente más pesada.

Key words: Soil gas CO₂, C isotopes, Seismicity, Rabaul, Papua New Guinea

Geogaceta, 20 (4) (1996), 1000-1003
ISSN: 0213683X

Introduction

Significant amounts of CO₂ are released to the atmosphere from active volcanic areas not only during eruptive stages but also during quiescent periods. This CO₂ emission from volcanic areas does not only occur from plumes and fumaroles, but also as diffuse outgassing from the surface environment (Aubert and Baubron, 1988; Allard *et al.*, 1991; Baubron *et al.*, 1990; Farrar *et al.*, 1995). Diffuse degassing monitoring in active volcanic systems is becoming an important tool for seismic-volcanic prediction studies because it can be carried out far away from active craters where volcanologists might experience a high level of risk. Changes in gas and volatile concentrations in the soils in volcanic areas have been used to detect changes in fluid flow resulting from seismic activity and magmatic intrusion (Varekamp and Buseck, 1984; Williams, 1985; Badalamenti *et al.*, 1988; Giammanco *et al.*, 1995; Farrar *et al.*, 1995). These observations can be a very important issue for volcanic scenarios such as caldera and rift zones where an eruption might build up new volcanic vents. In previous studies, the analysis and evaluation of soil gas and volatile distribution patterns have been quite

useful to detect areas of high heat flow which can become potential sites for new volcanic vents in active calderas (Williams, 1985; Pérez and Williams, 1990; Pérez, 1992; Salazar *et al.*, 1995). The aim of this study is to evaluate the existence of gas-flow through the seismic annulus structure defined during the unrest period of Rabaul caldera (Mori and McKee, 1987) by means of soil gas CO₂ measurements along a transect.

Sampling and analytical procedures

Soil gas samples were collected by using PVC pipes which were inserted in the ground at 40 cm. depth on November 1995. The upper part of the PVC pipes were closed by a cap with a septum to allow soil gas collection by using a syringe, while the lower part were open to allow soil gas to enter inside the PVC containers. Soil gas samples were collected after 5 days to allow soil gas atmosphere equilibration inside the PVC pipes.

A soil gas transect (A-A') of 43 sampling sites was carried out from the south of Matupit Island to the Northern part of the caldera between November 21-27, 1995 (Fig.1) following a selected space-sampling according to geological and geophysical characteristics of the

area. This geochemical profile was designed to intersect the seismic annulus defined by the 1980s seismic-uplift crisis at Rabaul caldera (Mori and McKee, 1987) and to be far away from the active volcanic vents to avoid interferences of degassing from volcanic vents. Soil gas samples were transferred to 10 cc. pre-evacuated vials and CO₂ measurements were carried out by Gas Chromatograph. Selected soil gas samples were also analyzed for their isotopic composition of carbon in both soil CO₂ and CH₄ by a GC-IRMS in the Laboratory for Earthquake Chemistry at the University of Tokyo.

Results and discussion

Forty soil gas samples were collected along the N-S trending transect on November 27, 1995, and analyzed for soil gas CO₂ concentrations and carbon isotopic composition of soil gas CO₂ and CH₄ during the first week of December, 1995. Soil gas CO₂ concentrations varied widely from 1045 ppm CO₂ (0.1% CO₂) to 196335 ppm CO₂ (19.6% CO₂). Spatial variations of soil gas CO₂ concentrations along the transect are shown in Figure 2, soil gas CO₂ levels peaked (up to 19.63%) along the seismically active fault zone (Mori and

McKee, 1987), near the old airport, at some distance from both Tavurvur and Vulcan which did erupt simultaneously on September 1994 (McKee *et al.*, 1994). These anomalously high levels of soil CO₂ are lower than those observed on the flank of Mammoth Mountain where soil gas CO₂ concentrations exceeded 30% (> 90% at several locations) and is being responsible for the killing of trees in the area (Farrar *et al.*, 1995). One significant difference between these two scenarios is that soil gas CO₂ measurements at Rabaul were designed to be relatively far away from active volcanic centers in the caldera while soil gas CO₂ survey at Long Valley was performed to evaluate diffuse «flank» emissions of magmatic CO₂ near by Mammoth Mountain, a large dacitic volcano. In the case of Rabaul caldera, seismic activity related to the unrest of this volcanic structure since the 1970s have clearly defined a seismic annulus which represents the actual caldera boundaries. Airport's area, near the old airfield, is the only subaerial location where soil gas surveys can intersect this seismic structure to evaluate diffuse gas-flow through this seismological-defined fault and without being affected for significant levels of degassing from active volcanic centers in the caldera. After evaluating all these observations, we might suggest that relatively high observed soil gas CO₂ concentrations at Airport represent a good geochemical indicator of gas-flow along this structure which can be defined as an active fault by geophysical and geochemical observations. Previous studies of soil gases at Rabaul caldera revealed relatively high ²²²Rn levels (>6x B and >9 x B, where B= background = 1.57 pCi/l) and relatively low Hg⁰ (<7.7 ppb, Background= 23.0ppb of Hg⁰) where the highest soil gas CO₂ were observed in the 1995 survey. This coupled observation of relatively high ²²²Rn and low Hg⁰ had been interpreted in other studies as active flush zones where a combination of mass and heat transport can account for this soil gas signature (Williams, 1985; Pérez, 1992). High soil gas CO₂ concentrations at Airport area could be also related to the effects of a higher thermal gradient along this active fault zone which might be also responsible for the potential release of Hg⁰ from the surface environment.

Other anomalously high soil gas CO₂ concentrations were seen at the Matupit causeway where an additional fault has been defined by geochemical (anomalous soil gas ²²²Rn and Hg⁰) and geophysical methods (Pérez, 1992; McKee, personal




Fig.1 Rabaul caldera map and detail of soil CO₂ transect. Volcanic centers post-caldera (solid circles) and pre-caldera (empty rectangles). RVO, location of the Rabaul Volcano Observatory. Caldera's topographic limits (dashed line) and old airport (dark rectangle).

communication) and near by Sulphur Creek area. Relatively low concentrations were observed at others sections of the transect, but the lowest values were detected at Matupit Island where a maximum uplift was observed in relation to the 1983-85 seismic-volcanic crisis at Rabaul caldera (McKee *et al.*, 1985). These low soil gas CO₂ concentrations observed at Matupit agree quite well with

the observed high soil pH values (> 8.01) in the area (Pérez and Williams, 1991, Pérez, 1992). Soil pH measurements related to other areas of this transect showed background levels (6.01 < pH < 8.00).

Carbon isotopic analysis of soil gas CO₂ and CH₄ in six selected samples (Fig.2) are listed in Table 1. $\delta^{13}\text{C}$ values of CO₂ ranged from -29.8‰ to -18.4‰




Fig.2 Soil gas CO₂ transect at Rabaul caldera along transect A-A'. Soil gas samples selected for carbon isotope measurements are indicated.

Fig. 2.-Transecto de CO₂ en lossuelos de la caldera de Rabaul siguiendo un perfil A-A'. Las muestras de gases en suelos analizados para isótopos de carbono son indicados




Fig.3 Relationship between soil gas CO₂ concentrations and $\delta^{13}\text{C}-\text{CO}_2$ from Rabaul caldera. Dashed lines represent two different trends.

Fig. 3.- Relación entre la concentración de CO₂ en los suelos $\delta^{13}\text{C}-\text{CO}_2$ en la caldera de Rabaul. Las líneas discontinuas representan dos tendencias diferentes

suggesting mainly a biogenic origin for the CO₂. The relationship between soil gas CO₂ concentrations and carbon isotopic signature of the CO₂ for these

six samples reflects a very interesting pattern (Fig.3). An increase of soil gas CO₂ is accompanied by an increase in the $\delta^{13}\text{C}$ values. This trend showed a good

linear regression with a correlation coefficient of 0.83, but this diagram suggest also that the data's distribution might follow two different regression lines (dashed lines in the Fig.3). After selecting the four samples with heavier isotopic signature, diagram of Abiogenic CO₂ vs $\delta^{13}\text{C}$ was constructed (Fig.4). All the data fit quite well according to the following linear regression equation

$$\delta^{13}\text{C}(\text{‰})-\text{CO}_2 = -23.176 + 0.24288 [\text{CO}_2] \quad (1)$$

which show a correlation coefficient of 1.00. Assuming that $\delta^{13}\text{C}$ values for magmatic CO₂ is around - 5.0 (‰), we can estimate what levels of soil gas CO₂ would be expected to be related to a significant diffuse magmatic degassing. This assumption might be reasonable because volcanic-hydrothermal discharges from Tavurvur volcano showed a carbon isotopic signature of - 5.1‰ for the CO₂. Following this equation, soil gas CO₂ concentrations around 75% can be linked to a clear magmatic origin. These estimates are in good agreement with soil gas CO₂ data from Long Valley where clear magmatic signatures of CO₂ were observed in soil gas samples with abiogenic CO₂ concentrations > 80%. On the contrary, soil gases collected inside of active craters with CO₂ concentrations around 30% showed already a clear magmatic signature (Allard *et al.*, 1991, Hernández *et al.*, 1996). These main differences are mainly related to the environment where diffuse degassing occurs. In the case of volcanic craters where there is not vegetation, biogenic CO₂ cannot be produce or be a significant source for the soil gas CO₂ reservoir. This case is totally opposite to that of diffuse degassing far away from active volcanic craters where biogenic CO₂ production can be relatively high and where deep-seated CO₂ can rise through permeable structures. This scenario might account for a mixing process between two components with different concentrations and isotopic compositions. Therefore, a mixing process between a biogenic signature and a minor volcanogenic contribution of CO₂ might also account for the observed carbon isotope signatures of the soil CO₂ at Rabaul caldera.

Conclusions

Relatively anomalous soil gas CO₂ up to 20% have been observed where high levels of seismic activity have been taking place since the unrest of Rabaul

caldera. Carbon isotopic composition of the soil gases suggest mainly a biogenic origin for the soil gas CO₂ and CH₄, but a small contribution of deep-seated CO₂ might contribute to the observed range of the soil gas CO₂ carbon isotopic composition. This anomalous soil CO₂ gas at the seismic annulus may reflect a significant gas-flow through this structure; therefore, soil gas CO₂ monitoring in the seismic annulus could be quite important for the volcanic-seismic surveillance of Rabaul caldera.

Acknowledgments

We are grateful to the Rabaul Volcano Observatory and Laboratory for Earthquake Chemistry at the University of Tokyo for their assistance and logistical support. We thank also Pedro A. Hernández, Youichi Shimoike, and Masanori Sato for their help at the laboratory. This research was supported by a grant of the EU-STF Programme in Japan (N.M.P.) and the Geological Survey of Papua New Guinea.

References

Aubert M. and Baubron J.-C. (1988). *J. Volcanol. & Geotherm. Res.*, 35, 217-225
 Allard P., Carbonelle J., Dajčević D., Le Bronce J., Morel P., Robe M.C., Maurenas J.M., Faivre-Pierret R., Martin D., Sabroux J.C., and Zettwoog P. (1991). *Nature*, 351, 387-391.
 Badalamenti B., Gurrieri S., Hauser S., Parello F., and Valenza M. (1988). *Rend. Soc. Ital. Min. Petrol.*, 43, 893-899.
 Baubron J.-C., Allard P., and Toutain J.P. (1990). *Nature*, 344, 51-53.
 Farrar C.D., Sorey M.L., Evans W.C., Howle J.F., Kerr B.D., Kennedy B.M., King C.Y., and Southon J.R. (1995). *Nature*, 376, 675-678.
 Giammanco S., Gurrieri S., and Valenza M. (1995). *Bull. Volcanol.*, 57, 52-60.
 Hernández P.A., Pérez N.M., Alvarez C.E., and Wakita H. (1996). *Geogaceta* (in press)
 McKee C.O., Lowenstein P.L., de Saint Ours P., Talai B., Itikari I., and Mori J.J. (1984). *Bull. Volcanol.*, 47, 397-411.
 McKee C.O., de Saint Ours P., and Talai B. (1994). *Bull. GVN*, 19, 2-5.
 Mori J. and McKee C.O. (1987). *Science*, 235, 193-195.
 Pérez N.M. and Williams S.N. (1990). *EOS Trans. Amer. Geophys. Union*, 71, 649
 Pérez N.M. and Williams S.N. (1991). *Proc.*




Fig. 4. Relationship between soil gas CO₂ concentrations and δ¹³C-CO₂ of samples which showed a heavier signature along the transect A-A' at Rabaul caldera. Correlation coefficient (1.000). Empty rectangle represent chemical and isotopic region values for volcanic-magmatic discharges at Rabaul caldera.

Fig. 4.- Relación entre la concentración de CO₂ de un origen no Biogenico y el δ¹³C-CO₂ en las muestras que indican una firma isotópica mas pesada a lo largo

Sample	CO ₂ (%)	δ ¹³ C-CO ₂ (‰ vs PDB)	δ ¹³ C-CH ₄ (‰ vs PDB)
3	0.39	- 29.8	-
36	6.00	- 21.7	- 70.0
43	15.12	- 19.5	- 70.3
17	19.63	- 18.4	- 68.3
34	10.73	- 20.6	- 66.2
29	3.62	- 26.9	- 70.9

Table 1.- Carbon isotopic composition of soil gas CO₂ and CH₄ from Rabaul caldera, Papua New Guinea

Int. Conf. on Active Volcanoes and Risk Mitigation, Napoli'91, Italy.
 Pérez N.M. (1992). *Ph.D. Dissertation*, Univ. La Laguna, 471 p.
 Salazar J.M., Pérez N.M., McKee C.O., de Saint Ours P., and Talai B. (1995). *EOS Trans. Amer. Geophys. Union*,
 Varekamp J.C. and Buseck P.R. (1984). *Geology*, 12, 29-47.
 Williams S.N. (1985). *Science*, 229, 551-553.