

Catalytic Functionalization of C-H Bonds of Azulene by Carbene/Nitrene Incorporation

Javier Carreras,^{a§} Yanay Popowski,^{b§} Ana Caballero,^{a,*} Elizabeth Amir,^{b,*} and Pedro J. Pérez^{a,*}

^aLaboratorio de Catálisis Homogénea, Unidad Asociada al CSIC, CIQSO-Centro de Investigación en Química Sostenible and Departamento de Química, Universidad de Huelva, 21007-Huelva, Spain

^bDepartment of Polymers and Plastics Engineering, Shenkar College, Ramat-Gan 5252626, Israel

ABSTRACT: The selective catalytic functionalization of the C-H bond of azulene upon incorporation of carbene or nitrene units with metal-based catalysts is described. Ethyl diazoacetate or ArI=NTs are employed as carbene or nitrene precursors, respectively. The azulene derivatives are subsequently employed as building blocks toward more complex structures with potential use as biodegradable materials.


INTRODUCTION

Since its chemical synthesis nearly eighty years ago,¹ azulene C₁₀H₈, has been in the center of extensive research.² Isomeric with naphthalene (a white crystalline solid), azulene displays a characteristic deep blue color and a large dipole moment of 1.08 D, due to an asymmetric π -polarization between the 5- and 7-membered rings. These characteristics make azulene an attractive building block for the construction of advanced functional materials for optoelectronic devices, light-emitting diodes (OLEDs) or organic field-effect transistors (OFETs), among others.^{3,4} Azulene is also gaining interest due to its reversible stimuli-responsive nature,⁵ antibacterial⁶ properties or some


potential medicinal and bioimaging applications⁷ as well.

Despite the interest in the availability of azulene-modified skeletons toward their use in the above-mentioned systems, there are only a few examples of the direct incorporation of substituents onto this bicyclic molecule.⁸ The functionalization of C-H bonds at the 5-member ring and, particularly, the formation of carbon-carbon bonds are relevant for subsequent uses of the modified azulene. Scheme 1 contains the representative examples of such metal-catalyzed processes, showing arylation (a),⁹ alkylation (b),¹⁰ alkenylation (c),¹¹ alkynylation (d),¹² propargylation (e)¹³ and carbonylation (f)¹⁴

Scheme 1. Reported metal-catalyzed methods for azulene C-H bond functionalization and C-C bond formation (a-f) and stoichiometric C-N bond generation (g)



Scheme 2. Azulene C-H bond functionalization and C-C and C-N bond formation by carbene or nitrene insertion




reactions. On the other hand, the direct amination of azulene C-H bonds has been scarcely reported, mainly following the stoichiometric vicarious nucleophilic substitution of hydrogen¹⁵ (Scheme 1g), photoinduced amina-

tions involving *N*-centered radicals¹⁶ or preparation of azo derivatives.¹⁷

For some applications such as biocompatible oligomers and polymers, the availability of sites that can undergo readily substitution patterns, such as esters or amines, is highly desirable. In this sense, a potential strategy toward that end could consist of the incorporation of CHCO_2Et or NTs units from $\text{N}_2=\text{CHCO}_2\text{Et}$ (ethyl diazoacetate, EDA) or $\text{PhI}=\text{NTs}$, serving as carbene or nitrene sources, respectively. At variance with the large number of examples on metal-catalyzed carbene transfer from EDA to a plethora of organic substrates,¹⁸ we have found no precedent with azulene. Only a single metal free protocol was described half a century ago, leading to monosubstitution at C_1 (Scheme 2).¹⁹ Additionally, the functionalization via metal-catalyzed nitrene transfer remains unreported. In this work, we describe the metal-catalyzed (Cu or Rh) reaction of azulene and diazoacetates yielding mono- (at C_1) or disubstituted (C_1 and C_3) derivatives, the latter being symmetric- or asymmetrically substituted (Scheme 2). These derivatives have been employed as starting materials toward the construction of novel azulene-dimeric structures with ester or amide linkages between the azulene units. The direct amination at C_1 upon copper-catalyzed nitrene transfer from $\text{ArI}=\text{NTs}$ is also reported, in the first example of this reaction with azulene as the substrate.

Scheme 3. Catalysts screened for the reaction of azulene with ethyl diazoacetate



RESULTS AND DISCUSSION

On the basis of our experience in the design of late transition metal-based catalysts for carbene transfer reactions from diazocompounds and concomitant functionalization of unsaturated substrates,^{20,21} we studied the reaction of azulene and EDA with a series of previously described catalysts for such transformations. Coinage metals complexes with either trispyrazolylborate or *N*-heterocyclic carbene ligands or well-known dirhodium carboxylate complexes were tested (Scheme 3). Most of the experi-

ments led to negative results, either by no EDA consumption or by formation of an untreatable mixture of compounds (see Figure S1 in the SI). However, the copper complex $\text{Tp}^{(\text{CF}_3)_2, \text{Br}}\text{Cu}(\text{NCCH}_3)$ ²² afforded a mixture of two compounds **1** and **2** derived from the incorporation of one or two CHCO_2Et units at the azulene skeleton at positions C_1 or C_1, C_3 , respectively (Table 1). Optimization of reac-

Table 1. Catalytic functionalization of azulene by carbene transfer from ethyl diazoacetate catalyzed by $\text{Tp}^{(\text{CF}_3)_2, \text{Br}}\text{Cu}(\text{NCCH}_3)$.^a

entry	cat:EDA:azulene ^b	mmol EDA	1 %	2 %
1	1:50:100	1.98	37	0
2	1:75:50	0.62	28	6
3	1:120:50	9.4	11	39

^aConditions: azulene was dissolved in CH_2Cl_2 along with $\text{Tp}^{(\text{CF}_3)_2, \text{Br}}\text{Cu}(\text{NCCH}_3)$ and a solution of EDA in the same solvent was added for 14 h with a syringe pump under inert atmosphere. See the Table S1 in the SI for experimental details. ^bMmol ratio.

tion conditions showed that the use of 1-2 mol% of the catalyst with respect to azulene and adjusting the EDA:azulene ratio allows the preferential formation of **1** or **2**. The reactions included in Table 1 provide a substantial amounts of the functionalized products: 0.73 mmol of

Table 2. Catalytic functionalization of azulene with ethyl 2-phenyldiazoacetate.^a


entry	catalyst	cat:PhEDA:azulene ^b	3 %	4 %
1	$\text{Tp}^{(\text{CF}_3)_2, \text{Br}}\text{Cu}(\text{L})$	1:50:100	21	0
2	$\text{Rh}_2(\text{OOC}(\text{CF}_3)_4)$	1:50:100	40	0
3	$\text{Tp}^{(\text{CF}_3)_2, \text{Br}}\text{Cu}(\text{L})$	1:100:50	32	6
4	$\text{Rh}_2(\text{OOC}(\text{CF}_3)_4)$	1:100:50	0	46

^aConditions employed as described Table 1. L = CH_3CN . See Table S2 in the SI for full experimental details. ^bMmol ratio.

1 (entry 1) or 1.5 mmol of **2** (entry 3, along with 0.43 of **1**). Both products and the unreacted azulene are readily separated upon column chromatography.

We next planned the use of a different diazo derivative such as ethyl diazo(phenyl)acetate (PhEDA). We examined the previously optimized conditions with $\text{Tp}^{(\text{CF}_3)_2, \text{Br}}\text{Cu}(\text{NCCH}_3)$, as well as $\text{Rh}_2(\text{OOCF}_3)_4$ and $\text{Rh}_2(\text{OAc})_4$. The rhodium complex bearing trifluoroacetate ligands improved the reactivity of the copper complex

Table 3. Catalytic functionalization of azulene by nitrene insertion from ArINTs derivatives catalyzed by $\text{Tp}^{\text{Br}_3}\text{Cu}(\text{NCCH}_3)$.^a




entry	R	cat:ArINTs:azulene ^b	mmol ArINTs	7%
1	H	1:50:100	0.20	15
2	OiPr	1:50:100	0.80	26
3	SO ₂ tBu	1:50:100	0.80	43
4	SO ₂ tBu	1:50:100	1.40	40

^aConditions: Azulene was dissolved in CH_2Cl_2 along with $\text{Tp}^{\text{Br}_3}\text{Cu}(\text{NCCH}_3)$ and a solution of ArINTs in the same solvent was added for 14 h with a syringe pump under inert atmosphere (PhINTs was added in one portion). See Table S₃ in the SI for full experimental details. ^bMmol ratio.

with PhEDA as the reactant (Table 2). Using the optimal set of reaction conditions described above, and $\text{Rh}_2(\text{OOCF}_3)_4$ as the catalyst, the appropriate tuning of the diazo-azulene ratio led to 40% of ethyl 2-(azulen-1-yl)-2-phenylacetate (3) (entry 2) or 46% of the disubstituted derivative 4, as a mixture of diastereomers (entry 4), both reactions performed at a 2 mmol scale of azulene.

We also studied the consecutive addition of two different diazo reagents. First, *tert*-butyl diazoacetate was reacted with azulene in the presence of $\text{Tp}^{(\text{CF}_3)_2, \text{Br}}\text{Cu}(\text{NCCH}_3)$ to obtain the monosubstituted derivative (5) in good yield

Scheme 4. Synthesis of asymmetrically disubstituted azulenes.




(Scheme 4). After isolation, this compound was subjected to a second C-H functionalization process using PhEDA and $\text{Rh}_2(\text{OOCF}_3)_4$ as catalyst, affording the unsymmetrical bis-substituted azulene 6 as the only product (Scheme 4).

Our study was extended to copper-catalyzed nitrene transfer reactions from *N*-tosyliminoiodanes derivatives (ArINTs) to azulene. Initial screening of the catalysts shown in Scheme 3 identified $\text{Tp}^{\text{Br}_3}\text{Cu}(\text{NCCH}_3)$ as the complex with higher activity on this transformation. Besides the reaction conditions (reaction time, reactants ratio, catalyst loading) we also tested different nitrene sources, moving from the parent $\text{PhI}=\text{NTs}$ ²³ to more soluble derivatives with substituents in the *ortho* position of the aromatic ring (Table 3).²⁴ In all cases, the *N*-tosyl-1-amino azulene (7) was obtained as the only product in the reaction. This compound has been characterized by NMR as well as by X-ray crystallography (see Figures S8 and S20 in the SI for complete description). Better yields in the reaction were achieved using *ortho*-sulfonyl- or *ortho*-isopropoxy substituted *N*-tosyliminoiodane derivatives compared with $\text{PhI}=\text{NTs}$. The solubility of these reagents in dichloromethane allowed us the slow addition of the reagent, reducing decomposition pathways. On the optimized conditions *ortho*-sulfonyl-*N*-tosyliminoiodane led to 40% of *N*-tosyl-1-amino azulene in 1.4 mmol scale, in the first example, to the best of our knowledge, of azulene C-H bond catalytic functionalization by nitrene transfer.

The potential of the functionalized azulene derivatives as building blocks for the preparation of degradable azulene-based materials was demonstrated by constructing azulene-dimeric structures. First, the mono-substituted azulenes 1 and 3 were readily hydrolyzed under basic


Scheme 5. Synthesis of azulene-dimeric structures.



conditions (Scheme 5) to obtain the corresponding 2-(azulen-1-yl)acetic acid (**8**) and 2-(azulen-1-yl)-2-phenylacetic acid (**9**), respectively, in high yields. Next, the azulene-dimeric compounds **10-12** were prepared in good yields through Steglich coupling reactions²⁵ treating **8-9** with either ethylene glycol or ethylenediamine in the presence of *N,N*-dicyclohexylcarbodiimide (DCC) as coupling agent and with catalytic amounts of 4-dimethylaminopyridine (DMAP) (Scheme 5). These previously unknown compounds were fully characterized by NMR and HRMS, and the solid structure of the blue amide **11** was also determined upon growing single crystals from cold dichloromethane solution.

We studied the reactivity of the disubstituted azulene derivatives **2** and **4** as well. In a similar fashion to the mono-substituted azulene acids, hydrolyses of the di-functionalized azulene esters **2** and **4** were carried out under basic conditions, affording the corresponding az-


Scheme 6. Derivatization of di-functionalized azulenes **2,4**.



ulene diacids **13** and **14** in high yields (Scheme 6). Diacid **13** exhibited poor solubility in dichloromethane and other common organic solvents precluding esterification or amidation reactions. Nevertheless, it could be successfully converted into the corresponding bis(acyl chloride) by treatment with thionyl chloride for 10 min. at room temperature. The intermediate, which was found to decompose upon standing, was subsequently reacted with 2-(tritylthio)ethan-1-amine to obtain the disubstituted amide derivative **15** in moderate yield (Scheme 6).

Finally, we also examined further functionalization of amino azulene **7**. Alkylation of the amine was performed with ethyl 2-bromoacetate leading to a protected glycine-azulene derivative **16** in good yield (Scheme 7), as an example of the synthesis of an azulene-containing α -aminoacid derivative.

Scheme 7. Derivatization of 1-amino azulene **7**.



CONCLUSIONS

In summary, we have developed the first metal-catalyzed systems for azulene C-H bond functionalization by carbene/nitrene transfer using diazocompounds or $\text{ArI}=\text{NTs}$, respectively. The process is regioselective toward the C_1 position. By adjusting the diazo-azulene ratio, both mono or disubstituted azulene (C_1 and C_3 positions) derivatives can be targeted with high selectivity and good yields. Furthermore, asymmetric disubstituted products are also accessible *via* sequential addition of two different diazo reagents. These novel azulene derivatives can be further converted into azulene-dimeric compounds bearing *O*- or *N*- based bridges. Further use of such compounds toward the preparation of (bio)degradable azulene-based materials is currently under development.

EXPERIMENTAL SECTION

General Information. Unless otherwise stated, reactions were carried out under nitrogen atmosphere with standard Schlenk techniques. Solvents were purchased from commercial sources or dried in a solvent purification system (MB SPS-800, MBRAUN). Analytical thin layer chromatography was carried out using TLC-aluminium sheets with 0.2 mm of silica gel (Merck 60 F254) and UV light as visualizing agent. Chromatography purifications were carried out using silica gel (0.035-0.070 mm, 60 Å). Organic solutions were concentrated under reduced pressure on a Büchi rotary evaporator. NMR spectra were recorded at 298 K on an Agilent 400 MR spectrometer. Chemical shift values for ^1H and ^{13}C are reported as δ values (ppm) relative to the deuterated solvent and coupling constants (*J*) in Hz. X-ray diffraction measurements were performed on an ApexDuo (Bruker-AXS) diffractometer system, using $\text{MoK}\alpha$ ($\lambda = 0.7107 \text{ \AA}$) radiation. The analyzed crystals were grown by slow evaporation of acetone solutions and freeze-cooled to ca. 110 K. Mass spectra were obtained on a SYNAPT - High Definition Mass Spectrometry or Orbitrap Elite mass spectrometer. Ionization methods: ESI (positive or negative) and APPI (atmospheric pressure photo ionization). All commercially available compounds were used as received. Catalysts have been prepared by described methodologies: $\text{Tp}^{\text{Br}_3}\text{Cu}(\text{NCCH}_3)$,²⁶ $\text{Tp}^{\text{Mes}}\text{Cu}(\text{THF})$,^{20a,27} $\text{Tp}^{(\text{CF}_3)_2\text{Br}}\text{Cu}(\text{NCCH}_3)$,²² $\text{Tp}^{\text{Br}_3}\text{Ag}(\text{THF})$,²⁸ $\text{Tp}^{(\text{CF}_3)_2\text{Br}}\text{Ag}(\text{THF})$,²² Rh_2OAc_4 ,²⁹ Rh_2TFA_4 ,³⁰ IPrAuCl ,³¹ IPrCuCl .³² *N*-Tosyliminoiodanes derivatives (ArINTs , $\text{Ar} = \text{C}_6\text{H}_5$,²³ 2- $\text{SO}_2t\text{Bu-C}_6\text{H}_4$,²⁴ 2-*i*PrO- C_6H_4 ²⁴) were synthesized by described methods.

Ethyl 2-(azulen-1-yl)acetate (**1**). Azulene (507 mg, 3.95 mmol, 2 eq.) and $\text{Tp}^{(\text{CF}_3)_2\text{Br}}\text{Cu}(\text{NCCH}_3)$ (38.1 mg, 0.04 mmol, 0.02 eq.) were mixed in dry CH_2Cl_2 (10 mL) under inert atmosphere. To this solution, EDA (0.23 mL, 1.98 mmol, 1 eq.) in dry CH_2Cl_2 (5 mL) was added during 14 hours *via* syringe pump. Then, the solvent was evaporated and the crude was purified by flash column chromatography (petroleum ether/AcOEt 100:0 to 95:5) to obtain the azulene derivative **1** as a blue oil (158 mg, 37%) and to recover 267 mg (53%) of azulene. Average of two runs. ^1H NMR (400 MHz, CDCl_3) δ 8.35 (d, $J = 9.7$ Hz, 1H, CH_{arom}), 8.31 (d, $J = 9.4$ Hz, 1H, CH_{arom}), 7.89 (d, $J = 3.8$ Hz, 1H, CH_{arom}), 7.60 (t, $J = 9.9$ Hz, 1H, CH_{arom}), 7.36 (d, $J = 3.8$ Hz, 1H, CH_{arom}), 7.18 (t, $J = 9.8$ Hz, 1H, CH_{arom}), 7.14 (t, $J = 9.7$ Hz, 1H, CH_{arom}), 4.15 (q, $J = 7.1$ Hz, 2H, $\text{CO}_2\text{CH}_2\text{CH}_3$), 4.10 (s, 2H, $\text{CH}_2\text{CO}_2\text{Et}$), 1.24 (t, $J = 7.1$ Hz, 3H, $\text{CO}_2\text{CH}_2\text{CH}_3$) ppm. $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, CDCl_3) δ 172.0 ($\underline{\text{C}}\text{O}_2\text{Et}$), 141.1 (C_{arom}), 138.1 (CH_{arom}), 137.8 (CH_{arom}), 136.9 (CH_{arom}), 136.5 (C_{arom}), 133.9 (CH_{arom}), 123.0 (CH_{arom}), 122.5 (CH_{arom}), 121.8 (C_{arom}), 117.0 (CH_{arom}), 61.0 ($\text{CO}_2\text{CH}_2\text{CH}_3$), 34.0 ($\underline{\text{C}}\text{H}_2\text{CO}_2\text{Et}$), 14.4 ($\text{CO}_2\text{CH}_2\text{CH}_3$) ppm. Resonance assignment was carried out using COSY and HSQC experiments. HRMS (EI-magnetic sector) m/z : $[\text{M}]^+$ Calcd for $\text{C}_{14}\text{H}_{14}\text{O}_2$ 214.0994; Found 214.0988.

Diethyl 2,2'-(azulene-1,3-diyl)diacetate (**2**). Azulene (500 mg, 3.90 mmol, 1 eq.) and $\text{Tp}^{(\text{CF}_3)_2\text{Br}}\text{Cu}(\text{NCCH}_3)$ (75.1 mg, 0.08 mmol, 0.02 eq.) were mixed in dry CH_2Cl_2 (15 mL) under inert atmosphere. To this solution, EDA (1.09 mL, 9.36 mmol, 2.4 eq.) in dry CH_2Cl_2 (7 mL) was added during 14 hours *via* syringe pump. Then, the solvent was evaporated and the crude was purified by flash column chromatography (petroleum ether/AcOEt 95:5 to 90:10) to obtain the azulene derivative **2** as a blue oil (456 mg, 39%) and 92 mg (11%) of azulene derivative **1**. ^1H NMR (400 MHz, CDCl_3): δ 8.28 (d, $J = 9.5$ Hz, 2H, CH_{arom}), 7.85 (s, 1H, CH_{arom}), 7.58 (t, $J = 9.8$ Hz, 1H, CH_{arom}), 7.13 (t, $J = 9.9$ Hz, 2H, CH_{arom}), 4.14 (q, $J = 7.2$ Hz, 4H, $\text{CO}_2\text{CH}_2\text{CH}_3$), 4.06 (s, 4H, $\text{CH}_2\text{CO}_2\text{Et}$), 1.24 (t, $J = 7.1$ Hz, 6H, $\text{CO}_2\text{CH}_2\text{CH}_3$) ppm. $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, CDCl_3): δ 171.9 ($\underline{\text{C}}\text{O}_2\text{Et}$), 139.2 (CH_{arom}), 138.1 (CH_{arom}), 137.3 (C_{arom}), 133.9 (CH_{arom}), 122.6 (CH_{arom}), 120.6 (C_{arom}), 61.0 ($\text{CO}_2\text{CH}_2\text{CH}_3$), 33.8 ($\underline{\text{C}}\text{H}_2\text{CO}_2\text{Et}$), 14.4 ($\text{CO}_2\text{CH}_2\text{CH}_3$) ppm. Resonance assignment was carried out using COSY and HSQC experiments. HRMS (EI-magnetic sector) m/z : $[\text{M}]^+$ Calcd for $\text{C}_{18}\text{H}_{26}\text{O}_4$ 300.1362; Found 300.1365.

Ethyl 2-(azulen-1-yl)-2-phenylacetate (**3**). Azulene (250 mg, 1.95 mmol, 2 eq.) and $\text{Rh}_2(\text{OOCFF}_3)_4$ (12.8 mg, 0.02 mmol, 0.02 eq.) were mixed in dry CH_2Cl_2 (5 mL) under inert atmosphere. To this solution, PhEDA (185 mg, 0.97 mmol,

1 eq.) in dry CH_2Cl_2 (5 mL) was added during 14 hours *via* syringe pump. Then, the solvent was evaporated and the crude was purified by flash column chromatography (petroleum ether/AcOEt 100:0 to 95:5) to obtain the azulene derivative **3** as a blue oil (115 mg, 41%) and to recover 132 mg (53%) of azulene. Average of two runs. ^1H NMR (400 MHz, CDCl_3) δ 8.36 (d, $J = 3.9$ Hz, 1H, CH_{arom}), 8.34 (d, $J = 3.4$ Hz, 1H, CH_{arom}), 8.06 (d, $J = 4.0$ Hz, 1H, CH_{arom}), 7.60 (t, $J = 9.9$ Hz, 1H, CH_{arom}), 7.42 (d, $J = 4.0$ Hz, 1H, CH_{arom}), 7.41 – 7.38 (m, 2H, CH_{arom}), 7.36 – 7.32 (m, 2H, CH_{arom}), 7.29 – 7.25 (m, 1H, CH_{arom}), 7.17 (t, $J = 9.7$ Hz, 2H, CH_{arom}), 5.73 (s, 1H, AzCHCO_2Et), 4.26 (AB of ABX_3 ; $J_{\text{AB}} = 10.8$ Hz, $J_{\text{X}} = 7.1$ Hz, 2H, $\text{CO}_2\text{CH}_2\text{CH}_3$), 1.29 (t, $J = 7.1$ Hz, 3H, $\text{CO}_2\text{CH}_2\text{CH}_3$) ppm. $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, CDCl_3) δ 173.0 ($\underline{\text{C}}\text{O}_2\text{Et}$), 141.2 (C_{arom}), 139.7 (C_{arom}), 137.8 (CH_{arom}), 137.3 (CH_{arom}), 137.1 (CH_{arom}), 135.9 (C_{arom}), 133.3 (CH_{arom}), 128.6 (CH_{arom}), 128.4 (CH_{arom}), 127.1 (CH_{arom}), 126.0 (C_{arom}), 123.2 (CH_{arom}), 122.7 (CH_{arom}), 117.2 (CH_{arom}), 61.3 ($\text{CO}_2\text{CH}_2\text{CH}_3$), 50.3 (AzCHCO_2Et), 14.3 ($\text{CO}_2\text{CH}_2\text{CH}_3$) ppm. Resonance assignment was carried out using COSY and HSQC experiments. HRMS (EI-magnetic sector) m/z : $[\text{M}]^+$ Calcd for $\text{C}_{20}\text{H}_{18}\text{O}_2$ 290.1307; Found 290.1309.

Diethyl 2,2'-(azulene-1,3-diyl)bis(2-phenylacetate) (**4**). Azulene (250 mg, 1.95 mmol, 1 eq.) and $\text{Rh}_2(\text{OOCFF}_3)_4$ (26.7 mg, 0.04 mmol, 0.02 eq.) were mixed in dry CH_2Cl_2 (5 mL) under inert atmosphere. To this solution, PhEDA (779 mg, 4.10 mmol, 2.1 eq.) in dry CH_2Cl_2 (5 mL) was added during 14 hours *via* syringe pump. Then, the solvent was evaporated and the crude was purified by flash column chromatography (petroleum ether/AcOEt 95:5 to 90:10) to obtain the azulene derivative **4** (1:1 mixture of diastereomers) as a blue solid (404 mg, 46%). ^1H NMR (400 MHz, CDCl_3) δ 8.28 (d, $J = 8.3$ Hz, 2H, CH_{arom}), 8.27 (d, $J = 8.4$ Hz, 2H, CH_{arom}), 8.17 (s, 1H, CH_{arom}), 8.14 (s, 1H, CH_{arom}), 7.55 (t, $J = 9.8$ Hz, 1H, CH_{arom}), 7.55 (t, $J = 9.7$ Hz, 1H, CH_{arom}), 7.37 – 7.28 (m, 16H, CH_{arom}), 7.27 – 7.21 (m, 4H, CH_{arom}), 7.11 (t, $J = 10.0$ Hz, 2H, CH_{arom}), 7.10 (t, $J = 10.0$ Hz, 2H, CH_{arom}), 5.68 (s, 4H, AzCHCO_2Et), 4.23 (q, $J = 7.1$ Hz, 4H, $\text{CO}_2\text{CH}_2\text{CH}_3$), 4.21 (q, $J = 7.1$ Hz, 4H, $\text{CO}_2\text{CH}_2\text{CH}_3$), 1.27 (t, $J = 7.1$ Hz, 6H, $\text{CO}_2\text{CH}_2\text{CH}_3$), 1.22 (t, $J = 7.1$ Hz, 6H, $\text{CO}_2\text{CH}_2\text{CH}_3$) ppm. $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, CDCl_3) δ 172.8 ($\underline{\text{C}}\text{O}_2\text{Et}$), 172.8 ($\underline{\text{C}}\text{O}_2\text{Et}$), 139.5 (C_{arom}), 138.2 (CH_{arom}), 138.2 (CH_{arom}), 137.8 (CH_{arom}), 137.7 (CH_{arom}), 136.9 (C_{arom}), 136.8 (C_{arom}), 133.7 (CH_{arom}), 133.6 (CH_{arom}), 128.6 (CH_{arom}), 128.6 (CH_{arom}), 128.5 (CH_{arom}), 128.5 (CH_{arom}), 127.1 (CH_{arom}), 127.1 (CH_{arom}), 125.0 (C_{arom}), 124.9 (C_{arom}), 123.0 (CH_{arom}), 61.3 ($\text{CO}_2\text{CH}_2\text{CH}_3$), 61.3 ($\text{CO}_2\text{CH}_2\text{CH}_3$), 50.5 (AzCHCO_2Et), 50.4 (AzCHCO_2Et), 14.3 ($\text{CO}_2\text{CH}_2\text{CH}_3$), 14.3 ($\text{CO}_2\text{CH}_2\text{CH}_3$) ppm. Resonance assignment was carried out using COSY and

HSQC experiments. HRMS (EI-magnetic sector) m/z: [M]⁺ Calcd for C₃₀H₂₈O₄ 452.1988; Found 452.1983.

Tert-butyl 2-(azulen-1-yl)acetate (**5**). Azulene (500 mg, 3.87 mmol, 2 eq.) and Tp^{(CF₃)₂Br}Cu(NCCH₃) (37 mg, 0.04 mmol, 0.02 eq.) were mixed in dry CH₂Cl₂ (10 mL) under inert atmosphere. To this solution, tBuEDA (268 μL, 1.93 mmol, 1 eq.) in dry CH₂Cl₂ (5 mL) was added during 14 hours *via* syringe pump. Then, the solvent was evaporated and the crude was purified by flash column chromatography (petroleum ether/AcOEt 100:0 to 95:5) to obtain the azulene derivative **5** as a blue oil (219 mg, 47%) and to recover 253 mg (51%) of azulene. ¹H NMR (400 MHz, CDCl₃) δ 8.38 (d, *J* = 9.7 Hz, 1H, CH_{arom}), 8.32 (d, *J* = 9.4 Hz, 1H, CH_{arom}), 7.95 (d, *J* = 3.9 Hz, 1H, CH_{arom}), 7.60 (t, *J* = 9.9 Hz, 1H, CH_{arom}), 7.41 (d, *J* = 3.8 Hz, 1H, CH_{arom}), 7.19 (t, *J* = 9.5 Hz, 1H, CH_{arom}), 7.14 (t, *J* = 9.5 Hz, 1H, CH_{arom}), 4.06 (s, 2H, CH₂CO₂tBu), 1.49 (s, 9H, CO₂C(CH₃)₃) ppm. ¹³C{¹H} NMR (100 MHz, CDCl₃) δ 171.2 (C=O₂tBu), 140.9 (C_{arom}), 138.0 (CH_{arom}), 137.6 (CH_{arom}), 136.7 (CH_{arom}), 136.4 (C_{arom}), 133.8 (CH_{arom}), 122.7 (CH_{arom}), 122.4 (C_{arom}), 122.2 (CH_{arom}), 116.9 (CH_{arom}), 80.7 (CO₂C(CH₃)₃), 35.2 (CH₂CO₂tBu), 28.1 (CO₂C(CH₃)₃) ppm. Resonance assignment was carried out using COSY and HSQC experiments. HRMS (EI-magnetic sector) m/z: [M]⁺ Calcd for C₁₆H₁₈O₂ 242.1307; Found 242.1322.

Ethyl 2-(3-(2-(tert-butoxy)-2-oxoethyl)azulen-1-yl)-2-phenylacetate (**6**). Azulene derivative **5** (219 mg, 0.90 mmol, 1 eq.) and Rh₂(OOCFCF₃)₄ (11.9 mg, 0.02 mmol, 0.02 eq.) were mixed in dry CH₂Cl₂ (3 mL) under inert atmosphere. To this solution, PhEDA (215 mg, 1.13 mmol, 1.25 eq.) in dry CH₂Cl₂ (3 mL) was added during 14 hours *via* syringe pump. Then, the solvent was evaporated and the crude was purified by flash column chromatography (petroleum ether/AcOEt 95:5 to 90:10) to obtain the azulene derivative **6** as a blue oil (163 mg, 45%). ¹H NMR (400 MHz, CDCl₃) δ 8.19 (d, *J* = 9.7 Hz, 1H, CH_{arom}), 8.15 (d, *J* = 9.7 Hz, 1H, CH_{arom}), 7.88 (s, 1H, CH_{arom}), 7.44 (t, *J* = 9.8 Hz, 1H, CH_{arom}), 7.29 – 7.23 (m, 2H, CH_{arom}), 7.23 – 7.16 (m, 2H, CH_{arom}), 7.16 – 7.10 (m, 1H, CH_{arom}), 7.02 (t, *J* = 9.5 Hz, 1H, CH_{arom}), 6.97 (t, *J* = 9.6 Hz, 1H, CH_{arom}), 5.56 (s, 1H, AzCHCO₂Et), 4.12 (q, *J* = 7.1 Hz, 2H, CO₂CH₂CH₃), 3.88 (s, 2H, CH₂CO₂tBu), 1.31 (s, 9H, CO₂C(CH₃)₃), 1.15 (t, *J* = 7.1 Hz, 3H, CO₂CH₂CH₃) ppm. ¹³C{¹H} NMR (100 MHz, CDCl₃) δ 173.0, 171.0 (C=O₂Et, C=O₂tBu), 139.6 (C_{arom}), 138.4 (CH_{arom}), 138.0 (CH_{arom}), 137.4 (C_{arom}), 136.6 (C_{arom}), 134.1 (CH_{arom}), 133.2 (CH_{arom}), 128.3 (CH_{arom}), 128.5 (CH_{arom}), 127.1 (CH_{arom}), 124.7 (C_{arom}), 122.6 (CH_{arom}), 122.6 (CH_{arom}), 121.5 (C_{arom}), 80.8 (CO₂C(CH₃)₃), 61.3 (CO₂CH₂CH₃), 50.1 (AzCHCO₂Et), 35.3 (CH₂CO₂tBu), 28.1 (CO₂C(CH₃)₃), 14.3 (CO₂CH₂CH₃)

ppm. Resonance assignment was carried out using COSY and HSQC experiments. HRMS (EI-magnetic sector) m/z: [M]⁺ Calcd for C₂₆H₂₈O₄ 404.1988; Found 404.1999.

N-(Azulen-1-yl)-4-methylbenzenesulfonamide (**7**). Azulene (358 mg, 2.80 mmol, 2 eq.) and Tp^{Br₃}Cu(NCCH₃) (29 mg, 0.03 mmol, 0.02 eq.) were mixed in dry CH₂Cl₂ (3 mL) under inert atmosphere. To this solution, *ortho*-sulfonyl-*N*-tosyliminoiodane (690 mg, 1.40 mmol, 1 eq.) in dry CH₂Cl₂ (12 mL) was added during 7 hours *via* syringe pump. Then, the solvent was evaporated and the crude was purified by flash column chromatography (petroleum ether/AcOEt 8:2) to obtain the azulene derivative **7** as a blue solid (166 mg, 40%) and to recover 207 mg (58%) of azulene. Mp: 145–146 °C. ¹H NMR (500 MHz, CD₃OD) δ 8.21 (dd, *J* = 9.4, 0.5 Hz, 1H, CH_{arom}), 8.08 (d, *J* = 9.4 Hz, 1H, CH_{arom}), 7.56 – 7.50 (m, 2H, CH_{arom}), 7.49 – 7.45 (m, 2H, CH_{arom}), 7.19 – 7.14 (m, 3H, CH_{arom}), 7.09 (t, *J* = 9.7 Hz, 1H, CH_{arom}), 6.98 (t, *J* = 9.8 Hz, 1H, CH_{arom}), 2.31 (s, 3H, SO₂C₆H₄CH₃) ppm. ¹³C{¹H} NMR (126 MHz, CD₃OD) δ 144.7 (C_{arom}), 140.0 (C_{arom}), 139.5 (CH_{arom}), 138.8 (CH_{arom}), 138.2 (C_{arom}), 134.7 (CH_{arom}), 134.5 (CH_{arom}), 133.7 (C_{arom}), 130.4 (CH_{arom}), 128.2 (CH_{arom}), 124.2 (CH_{arom}), 124.2 (C_{arom}), 123.5 (CH_{arom}), 116.2 (CH_{arom}), 21.4 (SO₂C₆H₄CH₃) ppm. Resonance assignment was carried out using COSY and HSQC experiments. HRMS (EI-ion trap) m/z: [M+Na]⁺ Calcd for C₁₇H₁₅NO₂SNa 320,0716; Found 320,0711. CCDC 1848245.

2-(Azulen-1-yl)acetic acid (**8**). Azulene derivative **1** (165 mg, 0.77 mmol) was dissolved in EtOH (15 mL), followed by the addition of KOH 1M (10 mL). The reaction was refluxed for 1.5 h, cooled to rt. and acidified with HCl 1M. Then, the crude product was extracted with EtOAc (2 x 20 mL), the combined extracts were washed repeatedly with H₂O, dried over MgSO₄ and the solvent was removed under reduced pressure. The blue oil obtained was used without further purification (136 mg, 95%). ¹H NMR (400 MHz, CDCl₃) δ 8.31 (d, *J* = 3.5 Hz, 1H), 8.29 (d, *J* = 3.7 Hz, 1H), 7.86 (d, *J* = 4.0 Hz, 1H), 7.60 (t, *J* = 9.6 Hz, 1H), 7.35 (d, *J* = 4.0 Hz, 1H), 7.18 (t, *J* = 10.0 Hz, 1H), 7.15 (t, *J* = 9.6 Hz, 1H), 4.12 (s, 2H) ppm. ¹³C{¹H} NMR (100 MHz, CDCl₃) δ 178.3, 141.1, 138.2, 137.9, 137.1, 136.6, 133.8, 123.2, 122.7, 120.6, 117.1, 33.6 ppm. HRMS (APPI-Q-TOF) m/z: [M-H]⁻ Calcd for C₁₂H₉O₂ 185.0603; Found 185.0611.

2-(Azulen-1-yl)-2-phenylacetic acid (**9**). Azulene derivative **9** was prepared correspondingly to **8** *via* the hydrolysis of **3** (138 mg, 0.47 mmol), yielding a blue oil (117 mg, 95%) which was used without further purification. ¹H NMR (400 MHz, CDCl₃) δ 8.31 (d, *J* = 9.6 Hz, 1H), 8.29 (d, *J* = 9.6 Hz, 1H), 8.00 (d, *J* = 4.0 Hz, 1H), 7.58 (t, *J* = 10 Hz, 1H), 7.39 – 7.27 (m, 6H), 7.26 – 7.21 (m, 1H), 7.15 (t, *J* = 9.6 Hz, 1H), 7.14 (t, *J* = 9.8 Hz, 1H), 5.70 (s, 1H) ppm. ¹³C{¹H} NMR (100

MHz, CDCl₃) δ 178.5, 141.3, 138.9, 138.0, 137.7, 137.4, 136.0, 133.3, 128.7, 128.6, 127.4, 125.0, 123.5, 123.0, 117.3, 50.1 ppm. HRMS (EI-magnetic sector) m/z: [M]⁺ Calcd for C₁₈H₁₄O₂ 262.0994; Found 262.0990.

Ethane-1,2-diyl bis(2-(azulen-1-yl)acetate) (**10**). Ethylene glycol (6.8 mg, 0.11 mmol), DCC (54.5 mg, 0.26 mmol) and DMAP (1.3 mg, 0.011 mmol) were added to a solution of **8** (41 mg, 0.22 mmol) in dry DCM (10 mL) and the reaction was stirred at rt. for 16 h under nitrogen atmosphere. Then, the reaction was filtered and the solvent was removed under reduced pressure. The crude was dissolved in DCM (2 mL) and filtered through a 0.22 μm syringe filter. The product was isolated as a dark blue oil by silica chromatography using 15% EtOAc in hexane (56 mg, 64%). ¹H NMR (400 MHz, CDCl₃) δ 8.30 (d, *J* = 9.4 Hz, 2H), 8.28 (d, *J* = 9.4 Hz, 2H), 7.84 (d, *J* = 3.6 Hz, 2H), 7.59 (t, *J* = 9.6 Hz, 2H), 7.35 (d, *J* = 3.6 Hz, 2H), 7.15 (t, *J* = 9.7 Hz, 2H), 7.14 (t, *J* = 9.7 Hz, 2H), 4.26 (s, 4H), 4.03 (s, 4H) ppm. ¹³C{¹H} NMR (100 MHz, CDCl₃) δ 171.6, 141.1, 138.1, 137.8, 137.0, 136.5, 133.8, 123.1, 122.6, 121.3, 117.1, 62.5, 33.7 ppm. HRMS (EI-magnetic sector) m/z: [M]⁺ Calcd for C₂₆H₂₂O₄ 398.1518; Found: 398.1544.

N,N'-(ethane-1,2-diyl)bis(2-(azulen-1-yl)acetamide) (**11**). Ethylenediamine (6.6 mg, 0.11 mmol), DCC (54.5 mg, 0.26 mmol) and DMAP (1.3 mg, 0.011 mmol) were added to a solution of **8** (41 mg, 0.22 mmol) in dry DCM (10 mL) and the reaction was stirred at rt. for 16 h under nitrogen atmosphere. Then, the reaction was filtered and the solvent was removed under reduced pressure. The crude was dissolved in DCM (2 mL) and filtered through a 0.22 μm syringe filter. The product was isolated as a dark blue solid on deactivated (5% H₂O) basic alumina using 1% MeOH in EtOAc (52 mg, 60%). Mp: 183–184 °C. ¹H NMR (400 MHz, CDCl₃) δ 8.33 (d, *J* = 9.6 Hz, 2H), 8.11 (d, *J* = 9.6 Hz, 2H), 7.71 (d, *J* = 4.0 Hz, 2H), 7.61 (t, *J* = 9.6 Hz, 2H), 7.37 (d, *J* = 4.0 Hz, 2H), 7.18 (t, *J* = 9.7 Hz, 2H), 7.13 (t, *J* = 9.8 Hz, 2H), 5.71 (brs, 2H), 3.88 (s, 4H), 3.11 (d, *J* = 2.4 Hz, 4H) ppm. ¹³C{¹H} NMR (100 MHz, CDCl₃) δ 172.6, 141.3, 138.5, 138.3, 137.3, 136.9, 133.9, 123.5, 123.0, 122.0, 117.4, 39.6, 35.9 ppm. HRMS (EI-magnetic sector) m/z: [M+H]⁺ Calcd for C₂₆H₂₅O₂N₂ 397.1911; Found: 397.1922. CCDC 1829375.

Ethane-1,2-diyl bis(2-(azulen-1-yl)-2-phenylacetate) (**12**). Azulene derivative **12** was prepared correspondingly to **10** using **9** (25 mg, 0.096 mmol), ethylene glycol (3 mg, 0.048 mmol), DCC (30 mg, 0.14 mmol) and DMAP (1.2 mg, 0.01 mmol) in CHCl₃ at reflux, yielding a blue solid (11 mg, 43%). Mp: 159–160 °C. ¹H NMR (400 MHz, CDCl₃) δ 8.29 (d, *J* = 9.4 Hz, 1H) 8.28 (d, *J* = 9.4 Hz, 1H), 8.24 (d, *J* = 6.2 Hz, 1H), 8.21 (d, *J* = 6.2 Hz, 1H), 7.94 (t, *J* = 4.0 Hz, 2H), 7.56 (t, *J* = 9.8 Hz, 1H), 7.55 (t, *J* = 9.8 Hz, 1H), 7.32–7.20 (m, 12H), 7.16–7.04 (m, 4H), 5.63 (s, 1H), 5.62 (s, 1H), 4.35 (s, 4H) ppm. ¹³C{¹H} NMR (100 MHz, CDCl₃) δ 172.8, 141.2,

139.3, 137.9, 137.3, 137.2, 135.9, 133.3, 128.6, 128.5, 127.2, 125.6, 123.3, 122.8, 117.3, 62.8, 50.1 ppm. HRMS (EI-magnetic sector) m/z: [M]⁺ Calcd for C₃₈H₃₀O₄ 550.2144; Found: 550.2143.

2,2'-(Azulene-1,3-diyl)diacetic acid (**13**). Diacid **13** was prepared correspondingly to **8** via the hydrolysis of **2** (150 mg, 0.5 mmol), yielding a blue solid (85 mg, 70%) which was used without further purification. Mp: 220–221 °C. ¹H NMR (400 MHz, DMSO-*d*₆) δ 12.33 (brs, 2H), 8.30 (d, *J* = 9.8 Hz, 2H), 7.77 (s, 1H), 7.63 (t, *J* = 9.8 Hz, 1H), 7.15 (t, *J* = 9.8 Hz, 2H), 4.00 (s, 4H) ppm. ¹³C{¹H} NMR (100 MHz, DMSO-*d*₆) δ 172.9, 139.1, 138.1, 136.5, 134.0, 122.0, 121.3, 33.0 ppm. HRMS (EI-magnetic sector) m/z: [M-H]⁻ Calcd for C₁₄H₁₁O₄ 243.0633; Found 243.0662.

2,2'-(Azulene-1,3-diyl)bis(2-phenylacetic acid) (**14**). Diacid **14** was prepared correspondingly to **8** via the hydrolysis of **4** (167 mg, 0.37 mmol), yielding a dark blue solid (139 mg, 95%, mixture of diastereomers) which was used without further purification. ¹H NMR (400 MHz, CDCl₃) δ 8.44 (s, 1H), 8.21 (d, *J* = 9.8 Hz, 2H), 8.19 (s, 1H), 8.12 (d, *J* = 9.8 Hz, 2H), 7.56 (t, *J* = 9.6 Hz, 1H), 7.51 (t, *J* = 9.6 Hz, 1H), 7.32–7.20 (m, 16H), 7.10 (t, *J* = 9.8 Hz, 2H), 7.03 (t, *J* = 9.8 Hz, 2H), 5.72 (s, 1H), 5.70 (s, 1H). ¹³C{¹H} NMR (100 MHz, CDCl₃) δ 180.1, 179.1, 138.9, 138.8, 138.7, 137.1, 136.2, 134.4, 133.8, 128.8, 128.7, 128.6, 127.5, 123.6, 123.5, 123.3, 50.9, 50.2 ppm. HRMS (EI-Q-TOF) m/z: [M+Na]⁺ Calcd for C₂₆H₂₀O₄Na 419,1254; Found: 419,1255.

2,2'-(Azulene-1,3-diyl)bis(N-(2-(tritylthio)ethyl)acetamide) (**15**). A flame-dried flask containing **13** (40 mg, 0.16 mmol) was cooled in an ice bath and SOCl₂ (230 μL, 3.2 mmol) was added dropwise under nitrogen atmosphere. Upon completion, the bath was removed and the reaction was stirred at rt. for 10 min., followed by removal of the SOCl₂ excess under reduced pressure. Then, the red-brown residue was dissolved in dry DCM (10 mL), 2-(tritylthio)ethan-1-amine (102 mg, 0.32 mmol) and Et₃N (180 μL, 1.28 mmol) were added and the reaction was stirred at rt. for 16 h under nitrogen atmosphere. After removal of the solvent under reduced pressure the product was isolated as a blue solid by silica chromatography using 80% EtOAc in hexane (9.7 mg, 20%). Mp: 202–203 °C. ¹H NMR (400 MHz, CDCl₃) δ 8.20 (d, *J* = 9.6 Hz, 2H), 7.69 (s, 1H), 7.51 (t, *J* = 9.6 Hz, 1H), 7.24–7.16 (m, 30H), 7.07 (t, *J* = 9.6 Hz, 2H), 5.46 (t, *J* = 5.6 Hz, 2H), 3.91 (s, 4H), 2.90 (q, *J* = 6.2 Hz, 4H), 2.27 (t, *J* = 6.4 Hz, 4H) ppm. ¹³C{¹H} NMR (100 MHz, CDCl₃) δ 171.2, 144.7, 139.6, 138.9, 137.8, 134.6, 129.5, 128.0, 126.9, 123.4, 121.3, 66.6, 38.1, 35.8, 31.9 ppm. HRMS (EI-magnetic sector) m/z: [M+Na]⁺ Calcd for C₅₆H₅₀O₂N₂S₂Na 869.3206; Found: 869.3217.

Ethyl N-(azulen-1-yl)-N-tosylglycinate (**16**). Amino azulene derivative **7** (44 mg, 0.15 mmol) and K₂CO₃ (41 mg, 0.30 mmol, 2 eq.) were dissolved in acetonitrile (2 mL), fol-

lowed by the addition of ethyl 2-bromoacetate (25 μ L, 0.22 mmol, 1.5 eq.), and the reaction was stirred at rt. for 5 hours. Then, the solvent was evaporated and the crude was purified by flash column chromatography (petroleum ether/Et₂O 1:1) to obtain the azulene derivative **16** as a blue oil (39 mg, 68%). ¹H NMR (500 MHz, CDCl₃) δ 8.63 (d, J = 9.7 Hz, 1H, CH_{arom}), 8.29 (d, J = 9.5 Hz, 1H, CH_{arom}), 7.67 (t, J = 9.7 Hz, 1H, CH_{arom}), 7.55 – 7.51 (m, 2H, CH_{arom}), 7.35 (d, J = 4.2 Hz, 1H, CH_{arom}), 7.31 – 7.20 (m, 4H, CH_{arom}), 7.16 (d, J = 4.2 Hz, 1H, CH_{arom}), 4.49 (s, 2H, NCH₂CO₂Et), 4.10 (q, J = 7.2 Hz, 2H, CO₂CH₂CH₃), 2.42 (s, 3H, SO₂C₆H₄CH₃), 1.17 (t, J = 7.2 Hz, 3H, CO₂CH₂CH₃) ppm. ¹³C{¹H} NMR (126 MHz, CDCl₃) δ 169.2 (C=O₂Et), 143.7 (C_{arom}), 139.1 (C_{arom}), 139.0 (CH_{arom}), 138.3 (CH_{arom}), 136.3 (C_{arom}), 135.8 (C_{arom}), 135.3 (CH_{arom}), 134.8 (CH_{arom}), 129.4 (CH_{arom}), 128.1 (CH_{arom}), 125.7 (C_{arom}), 124.4 (CH_{arom}), 115.1 (CH_{arom}), 61.5 (CO₂CH₂CH₃), 54.5 (NCH₂CO₂Et), 21.7 (SO₂C₆H₄CH₃), 14.2 (CO₂CH₂CH₃) ppm. Resonance assignment was carried out using COSY and HSQC experiments. HRMS (EI-ion trap) m/z : [M+H]⁺ Calcd for C₂₁H₂₂NO₄S 384.1264; Found 384.1259.

Additional compounds: *Benzyl 2-(azulen-1-yl)-2-phenylacetate* (**S1**). Benzyl alcohol (11 mg, 0.1 mmol), DCC (31 mg, 0.26 mmol) and DMAP (1.2 mg, 0.01 mmol) were added to a solution of **9** (26 mg, 0.1 mmol) in CHCl₃ (5 mL) and the reaction was refluxed for 16 h under nitrogen atmosphere. Then, the reaction was filtered and the solvent was removed under reduced pressure. The crude was dissolved in DCM (2 mL) and filtered through a 0.22 μ m syringe filter. The product was isolated as a dark blue oil by silica chromatography using 5% EtOAc in hexane (20 mg, 57%). ¹H NMR (400 MHz, CDCl₃) δ 8.31 (d, J = 9.8 Hz, 1H), 8.28 (d, J = 10.1 Hz, 1H), 7.99 (d, J = 3.6 Hz, 1H), 7.56 (t, J = 9.6 Hz, 1H), 7.37 (m, 3H), 7.31–7.23 (m, 8H), 7.14 (t, J = 9.7 Hz, 1H), 7.10 (t, J = 9.7 Hz, 1H), 5.76 (s, 1H), 5.21 (s, 2H) ppm. ¹³C{¹H} NMR (100 MHz, CDCl₃) δ 172.9, 141.2, 139.4, 137.8, 137.4, 137.1, 135.9, 133.3, 128.6, 128.6, 128.5, 128.3, 127.2, 125.8, 123.3, 122.7, 117.2, 67.0, 50.3 ppm. HRMS (EI-magnetic sector) m/z : [M]⁺ Calcd for C₂₅H₂₀O₂ 352.1463; Found: 352.1468.

Dibenzyl 2,2'-(azulene-1,3-diyl)bis(2-phenylacetate) (**S2**). Benzyl alcohol (14 mg, 0.13 mmol), DCC (34 mg, 0.16 mmol) and DMAP (1.6 mg, 0.013 mmol) were added to a solution of **14** (26 mg, 0.065 mmol) in CHCl₃ (5 mL) and the reaction was refluxed for 16 h under nitrogen atmosphere. Then, the reaction was filtered and the solvent was removed under reduced pressure. The crude was dissolved in DCM (2 mL) and filtered through a 0.22 μ m syringe filter. The product was isolated as a blue viscous oil (mixture of diastereomers) by silica chromatography using 5% EtOAc in hexane (23 mg, 61%). ¹H NMR (400 MHz, CDCl₃) δ 8.21 (d, J = 9.6 Hz, 1H), 8.20 (d, J = 9.6 Hz, 1H), 8.11 (s, 1H), 7.51 (t, J = 10.0 Hz, 1H), 7.29 – 7.21 (m, 20H), 7.03 (t, J = 9.6 Hz, 2H), 5.70 (s, 2H), 5.16 (s, 4H) ppm. ¹³C{¹H} NMR (100 MHz, CDCl₃) δ 172.7, 139.3, 139.2, 138.2, 137.9, 137.6, 136.9, 136.9, 135.9, 135.8, 133.8, 133.7,

128.6, 128.5, 128.2, 127.2, 124.6, 123.1, 67.1, 67.0, 50.4 ppm. HRMS (EI-magnetic sector) m/z : [M]⁺ Calcd for C₄₀H₃₂O₄ 576.2301; Found: 576.2294.

Supporting Information

NMR spectra (PDF) and cif files for compounds **7** and **11**.

AUTHOR INFORMATION

Corresponding Author

* perez@dqcm.uhu.es, ana.caballero@dqcm.uhu.es, eamir@shenkar.ac.il

Author Contributions

[§]These authors contributed equally.

ACKNOWLEDGMENT

We thank MINECO (CTQ2017-82893-C2-1-R and Juan de la Cierva fellowship for J.C.), Junta de Andalucía for Grant P12-FQM-1765 and CHAOS COST ACTION CA-15106. We also thank Manuel A. Pulido for preliminary experiments and Francisco Molina for X-ray structure determination.

REFERENCES

- (1) Pfau, A. St.; Plattner, P. A. Zur Kenntnis der flüchtigen Pflanzenstoffe VIII. Synthese des Vetivazulens. *Helv. Chim. Acta.* **1939**, *22*, 202-208.
- (2) (a) Preethalayam, P.; Krishnan, K. S.; Thulasi, S.; Chand, S. S.; Joseph, J.; Nair, V.; Jaroschik, F.; Radhakrishnan, K. V. Recent Advances in the Chemistry of Pentafulvenes. *Chem. Rev.* **2017**, *117*, 3930-3989. (b) Xin, H.; Gao, X. Application of Azulene in Constructing Organic Optoelectronic Materials: New Tricks for an Old Dog. *ChemPlusChem* **2017**, *82*, 945-956. (c) Skov, A. B.; Broman, S. L.; Gertsen, A. S.; Elm, J.; Jevric, M.; Cacciarini, M.; Kadziola, A.; Mikkelsen, K. V.; Nielsen, M. B. Aromaticity-Controlled Energy Storage Capacity of the Dihydroazulene-Vinylheptafulvene Photochromic System. *Chem. Eur. J.* **2016**, *22*, 14567-14575. (d) Gordon, M. The Azulenes. *Chem. Rev.* **1952**, *50*, 127-200.
- (3) (a) Nishimura, H.; Ishida, N.; Shimazaki, A.; Wakamiya, A.; Saeki, A.; Scott, L. T.; Murata, Y. Hole-Transporting Materials with a Two-Dimensionally Expanded π -System around an Azulene Core for Efficient Perovskite Solar Cells. *J. Am. Chem. Soc.* **2015**, *137*, 15656-15659. (b) Puodziukynaitė, E.; Wang, H.-W.; Lawrence, J.; Wise, A. J.; Russell, T. P.; Barnes, M. D.; Emrick, T. Azulene Methacrylate Polymers: Synthesis, Electronic Properties, and Solar Cell Fabrication. *J. Am. Chem. Soc.* **2014**, *136*, 11043-11049. (c) Yamaguchi, Y.; Ogawa, K.; Nakayama, K.; Ohba, Y.; Katagiri, H. Terazulene: A High-Performance n-Type Organic Field-Effect Transistor Based on Molecular Orbital Distribution Control. *J. Am. Chem. Soc.* **2013**, *135*, 19095-19098.
- (4) (a) López-Alled, L. M.; Sanchez-Fernandez, A.; Edler, K. J.; Sedgwick, A. C.; Bull, S. D.; McMullin, C. L.; Kociok-Köhn, G.;

- James, T. D.; Wenk, J.; Lewis, S. E. Azulene-boronate esters: colorimetric indicators for fluoride in drinking water. *Chem. Commun.* **2017**, 53, 12580-12583. (b) Xin, H.; Ge, C.; Yang, X.; Gao, H.; Yang, X.; Gao, X. Biazulene diimides: a new building block for organic electronic materials. *Chem. Sci.* **2016**, 7, 6701-6705. (c) Yamaguchi, Y.; Takubo, M.; Ogawa, K.; Nakayama, K.-I.; Koganezawa, T.; Katagiri, H. Terazulene Isomers: Polarity Change of OFETs through Molecular Orbital Distribution Contrast. *J. Am. Chem. Soc.* **2016**, 138, 11335-11343. (d) Barman, S.; Furukawa, H.; Blacque, O.; Venkatesan, K.; Yaghi, O. M.; Berke, H. Azulene based metal-organic frameworks for strong adsorption of H₂. *Chem. Commun.* **2010**, 46, 7981-7983.
- (5) Amir, E.; Amir, R. J.; Campos, L. M.; Hawker, C. J. Stimuli-Responsive Azulene-Based Conjugated Oligomers with Polyani-line-like Properties. *J. Am. Chem. Soc.* **2011**, 133, 10046-10053.
- (6) López-Pérez, B.; Pepper, H. P.; Ma, R.; Fawcett, B. J.; Peheré, A. D.; Wei, Q.; Ji, Z. C.; Polyak, S. W.; Dai, H. Q.; Song, F. H.; Abell, A. D.; Zhang, L. X.; George, J. H. Biosynthetically Guided Structure-Activity Relationship Studies of Merochlorin A, an Antibiotic Marine Natural Product. *ChemMedChem* **2017**, 12, 1969-1976.
- (7) Zhou, Y.; Zhuang, Y.; Li, Xin; Ågren, H.; Yu, L.; Ding, J.; Zhu, L. Selective Dual-Channel Imaging on Cyanostyryl-Modified Azulene Systems with Unimolecularly Tunable Visible-Near Infrared Luminescence. *Chem. Eur. J.* **2017**, 23, 7642-7647.
- (8) Shi, X.; Sasmal, A.; Soulé, J.-F.; Doucet, H. Metal-Catalyzed C-H Bond Activation of 5-Membered Carbocyclic Rings: A Powerful Access to Azulene, Acenaphthylene and Fulvene Derivatives. *Chem. Asian J.* **2018**, 13, 143-157.
- (9) (a) Dyker, G.; Borowski, S.; Heiermann, J.; Körning, J.; Opwis, K.; Henkel, G.; Köckerling, M. First intermolecular palladium catalyzed arylation of an unfunctionalized aromatic hydrocarbon. *J. Organomet. Chem.* **2000**, 606, 108-111. (b) Murai, M.; Yanagawa, M.; Nakamura, M.; Takai, K. Palladium-Catalyzed Direct Arylation of Azulene Based on Regioselective C-H Bond Activation. *Asian J. Org. Chem.* **2016**, 5, 629-635. (c) Cambeiro, X. C.; Ahlsten, N.; Larrosa, I. Au-Catalyzed Cross-Coupling of Arenes via Double C-H Activation. *J. Am. Chem. Soc.* **2015**, 137, 15636-15639.
- (10) (a) Dyker, G.; Muth, E.; Hashmi, A. S. K.; Ding, L. Gold(III) Chloride-Catalyzed Addition Reactions of Electron-Rich Arenes to Methyl Vinyl Ketone. *Adv. Synth. Catal.* **2003**, 345, 1247-1252. (b) Aguilar, D.; Contel, M.; Navarro, R.; Urriolabeitia, E. P. Organogold(III) Iminophosphorane Complexes as Efficient Catalysts in the Addition of 2-Methylfuran and Electron-Rich Arenes to Methyl Vinyl Ketone. *Organometallics* **2007**, 26, 4604-4611. (c) Timmerman, J. C.; Schmitt, W. W.; Widenhoefer, R. A. Gold-Catalyzed Intermolecular, anti-Markovnikov Hydroarylation of Methylene-cyclopropanes with Indoles. *Org. Lett.* **2016**, 18, 4966-4969. (d) Gu, L. Wolf, L. M. Zielinski, A. Thiel, W. Alcarazo, M. α -Dicationic Chelating Phosphines: Synthesis and Application to the Hydroarylation of Dienes. *J. Am. Chem. Soc.* **2017**, 139, 4948-4953.
- (11) Park, S.; Yong, W.-S.; Kim, S.; Lee, P. H. Diastereoselective N-Sulfonylaminoalkenylation of Azulenes from Terminal Alkynes and Azides via N-Sulfonyl-1,2,3-triazoles. *Org. Lett.* **2014**, 16, 4468-4471.
- (12) Székely, A.; Peter, A.; Aradi, K.; Tolnai, G. L.; Novak, Z. Gold-Catalyzed Direct Alkynylation of Azulenes. *Org. Lett.* **2017**, 19, 954-957.
- (13) (a) Nishibayashi, Y.; Yoshikawa, M.; Inada, Y.; Hidai, M.; Uemura, S. Ruthenium-Catalyzed Propargylation of Aromatic Compounds with Propargylic Alcohols. *J. Am. Chem. Soc.* **2002**, 124, 11846-11847. (b) Inada, Y.; Yoshikawa, M.; Milton, M. D.; Nishibayashi, Y.; Uemura, S. Ruthenium-Catalyzed Propargylation of Aromatic Compounds with Propargylic Alcohols. *Eur. J. Org. Chem.* **2006**, 881-890.
- (14) Liu, J.; Li, H.; Dühren, R.; Liu, J.; Spannenberg, A.; Franke, R.; Jackstell, R.; Beller, M. Markovnikov-Selective Palladium Catalyst for Carbonylation of Alkynes with Heteroarenes. *Angew. Chem. Int. Ed.* **2017**, 56, 11976-11980.
- (15) See for example: Makosza, M.; Podraza, R. Hydroxylation and Amination of Azulenes by Vicarious Nucleophilic Substitution of Hydrogen. *Eur. J. Org. Chem.* **2000**, 193-198.
- (16) (a) Davies, J.; Svejstrup, D.; Fernandez-Reina, D.; Sheikh, N. S.; Leonori, D. Visible-Light-Mediated Synthesis of Amidyl Radicals: Transition-Metal-Free Hydroamination and N-Arylation Reactions. *J. Am. Chem. Soc.* **2016**, 138, 8092-8095. (b) Svejstrup, T. D.; Ruffoni, A.; Juli, F.; Aubert, V. M.; Leonori, D. Synthesis of Arylamines via Aminium Radicals. *Angew. Chem. Int. Ed.* **2017**, 56, 14948-14952.
- (17) (a) Lacroix, P. G.; Malfant, I.; Iftime, G.; Razus, A. C.; Nakatani, K.; Delaire, J. A. Azo-Azulene Derivatives as Second-Order Nonlinear Optical Chromophores. *Chem. Eur. J.* **2000**, 6, 2599-2608. (b) Tskhovrebov, A. G.; Naested, L. C. E.; Solari, E.; Scopelliti, R.; Severin, K. Synthesis of Azoimidazolium Dyes with Nitrous Oxide. *Angew. Chem. Int. Ed.* **2015**, 54, 1289-1292.
- (18) (a) Cheng, Q.-Q.; Doyle, M. P. The Selection of Catalysts for Metal Carbene Transformations. *Adv. Organomet. Chem.* **2016**, 66, 1-31. (b) Doyle, M. P.; McKervey, M. A.; Ye, T. *Modern Catalytic Methods for Organic Synthesis with Diazo Compounds*, John Wiley & Sons, New York, 1998;
- (19) Anderson, A. G.; Rhodes, R. C. The Reaction of Azulenes with Aliphatic Diazo Compounds. *J. Org. Chem.* **1965**, 30, 1616-1619.
- (20) (a) Díaz-Requejo, M. M.; Caballero, A.; Belderrain, T. R.; Trofimenko, S.; Pérez, P. J. Copper(I)-Homoscorpionate Catalysts for the Preferential, Kinetically Controlled Cis Cyclopropanation of α -Olefins with Ethyl Diazoacetate. *J. Am. Chem. Soc.* **2002**, 124, 978-983. (b) Morilla, M. E.; Díaz-Requejo, M. M.; Belderrain, T. R.; Nicasio, M. C.; Trofimenko, S.; Pérez, P. J. Reaction of Ethyl Diazoacetate with Alkyl-Aromatic Substrates: Influence of the TpxCu Catalyst in the Addition versus Insertion Chemoselectivity (Tp^x = Homoscorpionate). *Organometallics* **2004**, 23, 293-295.
- (21) (a) Fructos, M. R.; Belderrain, T. R.; Frémont, P.; Scott, N. M.; Nolan, S. P.; Díaz-Requejo, M. M.; Pérez, P. J. A Gold Catalyst for Carbene-Transfer Reactions from Ethyl Diazoacetate. *Angew. Chem. Int. Ed.* **2005**, 44, 5284-5288. (b) Rivilla, I.; Gómez-Emeterio, B. P.; Fructos, M. R.; Díaz-Requejo, M. M.; Pérez, P. J. Exclusive Aromatic vs Aliphatic C-H Bond Functionalization by Carbene Insertion with Gold-Based Catalysts. *Organometallics* **2011**, 30, 2855-2860. (c) Rivilla, I.; Díaz-Requejo, M. M.; Pérez, P. J. Gold-catalyzed naphthalene functionalization. *Beilstein J. Org. Chem.* **2011**, 7, 653-657.
- (22) Gava, R.; Olmos, A.; Noverges, B.; Varea, T.; Álvarez, E.; Belderrain, T. R.; Caballero, A.; Asensio, G.; Pérez, P. J. Discovering Copper for Methane C-H Bond Functionalization. *ACS Catal.* **2015**, 5, 3726-3730.

(23) Yamada, Y.; Yamamoto, T.; Okawara, M. Synthesis and reaction of new type I-N ylide, *N*-tosyliminoiodinane. *Chem. Lett.* **1975**, *4*, 361-362.

(24) (a) Macikenas, D.; Skrzypczak-Jankun, E.; Protasiewicz J. D. A New Class of Iodonium Ylides Engineered as Soluble Primary Oxo and Nitrene Sources. *J. Am. Chem. Soc.* **1999**, *121*, 7164-7165. (b) Guo, M.; Dong, H.; Li, J.; Cheng, B.; Huang, Y.-Q.; Feng, Y.-Q.; Lei, A. Spectroscopic observation of iodopyrene metalloporphyrin adducts and manganese(V)-oxo porphyrin species in a cytochrome P450 analogue. *Nat. Commun.* **2012**, *3*, 1190-1198. (c) Yoshimura, A.; Nemykin, V. N.; Zhdankin, V. V. *o*-Alkoxyphenyliminoiodanes: Highly Efficient Reagents for the Catalytic Aziridination of Alkenes and the Metal-Free Amination of Organic Substrates. *Chem. Eur. J.* **2011**, *17*, 10538-10541.

(25) Neises, B.; Steglich, W. Simple Method for the Esterification of Carboxylic Acids. *Angew. Chem. Int. Ed.* **1978**, *17*, 522-524.

(26) Caballero, A.; Díaz-Requejo, M. M.; Belderráin, T. R.; Nicasio, M. C.; Trofimenko, S.; Pérez, P. J. Functionalization of Carbon-Hydrogen Bonds of Hydrocarbons and Ethers via Carbene Insertion with Copper(I)-Homoscorpionate Catalysts. *Organometallics* **2003**, *22*, 4145-4150. (b) Caballero, A.; Díaz-Requejo, M. M.; Belderráin, T. R.; Nicasio, M. C.; Trofimenko, S.; Pérez, P. J. Highly Regioselective Functionalization of Aliphatic Carbon-Hydrogen Bonds with a Perbromohomoscorpionate Copper(I) Catalyst. *J. Am. Chem. Soc.* **2003**, *125*, 1446-1447. (c) See

also: Trofimenko, S. *Scorpionates, The Coordination Chemistry of Polypyrazolylborate Ligands*; Imperial College Press: London, 1999.

(27) Díaz-Requejo, M. M.; Belderráin, T. R.; Nicasio, M. C.; Trofimenko, S.; Pérez, P. J. Intermolecular Copper-Catalyzed Carbon-Hydrogen Bond Activation via Carbene Insertion. *J. Am. Chem. Soc.* **2002**, *124*, 896-897.

(28) Urbano, J.; Belderráin, T. R.; Nicasio, M. C.; Trofimenko, S.; Díaz-Requejo, M. M.; Pérez, P. J. Functionalization of Primary Carbon-Hydrogen Bonds of Alkanes by Carbene Insertion with a Silver-Based Catalyst. *Organometallics* **2005**, *24*, 1528-1532.

(29) Rempel, G. A.; Legzdins, P.; Smith, H.; Wilkinson, G. Tetrakis(acetato)dirhodium(II) and Similar Carboxylato Compounds. *Inorg. Synth.* **1972**, *13*, 90-91.

(30) Telser, J.; Drago, R. S. Reactions of rhodium trifluoroacetate with various Lewis bases. Formation of 4:1 complexes with pyridine and tert-butyl isocyanide and bond cleavage with phosphorus donors. *Inorg. Chem.* **1984**, *23*, 2599-2606.

(31) Visbal, R.; Laguna, A.; Gimeno, M. C. Simple and efficient synthesis of [MCl(NHC)] (M = Au, Ag) complexes. *Chem. Commun.* **2013**, *49*, 5642-5644.

(32) Santoro, O.; Collado, A.; Slawin, A. M. Z.; Nolan, S. P.; Cazin, C. S. J. A general synthetic route to [Cu(X)(NHC)] (NHC = N-heterocyclic carbene, X = Cl, Br, I) complexes. *Chem. Commun.* **2013**, *49*, 10483-10485.

