

Trophic diversity of the otter (*Lutra lutra* L.) in temperate and Mediterranean freshwater habitats

5 **Miguel Clavero^{1,2}, José Prenda¹ and Miguel Delibes²**

¹ Departamento de Biología Ambiental y Salud Pública, Universidad de Huelva. Campus Universitario de El Carmen, Avd/ Fuerzas Armadas s/n. 21007 Huelva, Spain. E-mail: jprenda@uhu.es

² Departamento de Biología Aplicada. Estación Biológica de Doñana. CSIC. Pabellón del Perú, Avda. María Luisa s/n, 41013 Sevilla, Spain. E-mail: mdelibes@ebd.csic.es

Short title: Otter diet diversity in Europe

Keywords: feeding habits, freshwater ecosystems, freshwater fishes, geographic variation, otter, *Lutra lutra*, Mediterranean climate.

ABSTRACT

Aim: To analyse the geographical patterns in the composition and diversity of otter's (*Lutra lutra*, L.) diet and their relationship with climatic characteristics.

Location: European freshwater habitats under Mediterranean and temperate
5 climatic regimes.

Methods: 37 otter diet studies were reviewed, 21 from temperate and 16 from Mediterranean areas. All studies were based in spraint analysis and their results expressed as relative frequency of occurrence (RFO) of seven main prey categories. Principal Component Analysis (PCA) was performed to extract the main gradients of
10 diet composition. Pearson's correlation and T-tests were used to assess the relation between diet characteristics (composition, diversity, taxonomic richness) and geographic and climatic variables.

Results: A clear latitudinal gradient in trophic diversity and diet composition is observed. Otter diet is more diverse and features more prey classes in southern
15 localities, while the species is more piscivorous towards the north, where it predate upon a higher number of fish families. This pattern is contrasting when temperate and Mediterranean localities of Europe are compared. Mediterranean otters behave as more generalist predators than temperate ones, relying less on fish, and more on crayfish, aquatic invertebrates and reptiles.

Main conclusions: Geographical differences in otters feeding ecology in Europe
20 seem to be related with the two contrasted climatic conditions affecting prey populations. The otter can act as a highly specialised piscivorous predator in temperate freshwater ecosystems, which do not suffer a dry season and are

comparatively more stable than Mediterranean ones. However, the unpredictable prey availability in Mediterranean areas, affected by strong spatial and temporal water shortages, would favour a diversification of otter diet.

INTRODUCTION

The latitudinal gradient in diversity (i.e. a growing richness of flora and fauna species from the poles to the tropics) is considered the oldest (Hawkins 2001) and the best known (Rosenzweig 1995) biogeographical “pattern” of ecology. Since
5 abundance and diversity of potential prey are the most important features determining a predator’s niche width (MacArthur & Pianka, 1966; Schoener, 1971), we could expect, on a large scale, that predator species occupying a broad geographical range may experience variations of their food-niche breadth following changes in prey communities. At the intraspecific level, such differential use of
10 trophic resources among allopatric populations may occur with minor or no apparent morphological or physiological changes (Futuyma & Moreno, 1988; Martín *et al.*, 1995).

Studies on the trophic ecology of several species of birds and mammals support the prediction of a decreasing trophic diversity at higher latitudes. A classical
15 example is the puma (*Puma concolor*, L.), whose range covers from Alaska in the north to Tierra del Fuego in the south and whose highest food-niche breadths are reached in tropical America (Iriarte *et al.* 1990). In Europe, an increasing dietary diversity towards the south has been suggested for barn owl (*Tyto alba*, Scop.; Herrera, 1974), Montagu’s harrier (*Circus pigargus*, L.; Arroyo, 1997) and other
20 raptors (Korpimaki & Marti, 1995), among the birds, and common genet (*Genetta genetta*, L.; Virgós *et al.*, 1999) among the mammals.

Since abiotic environmental factors, such as climate regime, are in the last term the determinants of the composition and structure of prey communities (Smith &

Smith, 2000), the observed dietary patterns in Europe could be associated to the two principal climatic areas defined in the continent: Temperate in central and part of northern Europe and Mediterranean in the south (Fig. 1). Temperate climate is characterized by cold winters and mild and rainy summers, without a summer drought. Mediterranean areas (the term *Mediterranean* is always used in the text in a climatic, an not strictly geographic, sense), considered transitional between cold temperate and dry tropical zones, are characterized by the unique combination of hot dry summers and cool humid winters. Little or no surface water is available during the summer, generating a period of effective physiological drought. Added to this seasonality is a characteristic unpredictability, with dramatic variations of weather conditions among years, between seasons of a given year and even in the course of a single day (Blondel & Aronson, 1999).

Freshwater ecosystems are specially sensitive to predictable and unpredictable environmental variations such as those characteristic of Mediterranean climates (Prenda & Gallardo 1996, Gasith & Resh 1999, Magalhães *et al.*, 2002).

Consequently, the Eurasian otter (*Lutra lutra*, L.), a top predator of aquatic ecosystems, could experiment changes in its food and foraging behaviour according to the composition and structure of prey communities in Mediterranean and temperate ecosystems. The strong decline suffered by the populations of the otter in Europe from the 1950s has largely stimulated research into the species distribution and ecology (Mason & Macdonald, 1986), being the diet one of the central issues (Carrs, 1995). Such a situation provides a good chance to analyse the possible variations in otter's diet diversity throughout Europe.

Mason & Macdonald (1986) suggested that otters in southern Europe featured more amphibians and reptiles in their diets than those of the north. Also, Adrián & Delibes (1987) indicated that frequency of occurrence of insects, amphibians and reptiles in faeces of otters in Europe seemed to increase as latitude decreased. Again, 5 Ruiz-Olmo (1995) showed that reptiles were a common food of Mediterranean otters, but were very rare in other latitudes. However, a recent analysis of the diet of otters in Eurasia failed to detect any of these trends (Jedrzejewska *et al.*, 2001). These authors stated that otter diets in Eurasia do not change with latitude, but change with habitat: fish are more frequent as prey on sea shores, followed by lakes, 10 and rivers and streams; amphibians and crustaceans show just the contrary trend.

The aim of this paper is to search for geographical patterns in diet composition and diversity of otters living in Mediterranean and temperate climatic conditions of Europe. To remove the important bias that coastal otters can introduce, we limited our sources to freshwater habitats.

15 MATERIAL AND METHODS

FIGURE 1

Data on otter diet were taken from 37 diet studies from the available literature. Each study area was assigned to Mediterranean or temperate climate conditions according to the map of Emberger *et al.* (1963). This resulted in 16 Mediterranean and 21 temperate localities (Fig. 1). All the studies were based on spraint analysis 20 and the data expressed as relative frequency of occurrence (RFO) (number of occurrences of a certain item as percentage of the total number of occurrences of all prey items). Only studies with more than 200 occurrences of the different prey

categories were considered. Usually, results from several places or streams from the same area were pooled to avoid pseudoreplication (Hulbert 1994). Whenever necessary, original data were transformed to RFO to allow a correct comparison. Recent critical analyses (Carrs & Parkinson, 1996; Jacobsen & Hansen, 1996) have shown that RFO is not the best method to assess otter diet, leading to overestimation of medium size prey items and underestimation of the smaller and bigger ones. However, Jacobsen & Hansen (1996) compared several methods and found that similarity of RFO results with those of more accurate methods was between 80% and 90% (Renkonens Index of Similarity). Thus, being the most frequent method used in literature, and for our aim of comparisons, we consider RFO an appropriate methodology to establish dietary geographical patterns.

Seven basic prey categories were considered: fish, amphibians, reptiles, birds, mammals, crayfish and other aquatic invertebrates (beetles, damselfly nymphs, small shrimps, etc). For each local diet, we calculated the number of prey categories (NPC) and, when possible (in 35 studies), the total number of fish families (NFF) present in the sample. To quantify the general diet diversity for each location, the Shannon-Wiener expression (H') applied to frequencies of occurrence was used.

Principal Component Analysis (PCA) was performed to an arcsine transformed matrix of RFO x study locations ($n=37$) in order to summarise general patterns in otter diet throughout the study area. RFO were arcsine transformed prior the analysis to homogenise variances (Zar, 1984). Pearson's correlations between occurrence data, trophic and geographic variables and principal components were calculated. To assess differences between Mediterranean and temperate diets we used t-tests.

Whenever multiple t-test were performed, significance levels were corrected using the Bonferroni method (Sokal & Rohlf, 1995).

RESULTS

5 Fish was the otter's main food category, representing almost 75% of the consumed preys among otters inhabiting freshwater habitats (Table 1). The remaining categories can be considered as secondary in a general approximation to otter diet. Predation upon amphibians, crayfish and other invertebrates was frequent, while birds, reptiles and mammal were rather rare in otter diet.

10 The first principal component (PC 1) produced a strong ordination of the different study locations, explaining 40% of the total observed variance. PC 1 defined a gradient running from high and almost exclusive fish consumption to relative high predation upon aquatic invertebrates, amphibians, crayfish and reptiles (Table 1). The diet gradient represented by PC 1 showed a marked geographical component, being highly correlated with latitude ($r= 0.63$; $P < 0.001$) (Fig. 2A). The second principal component explained only 17.6% of the observed variance and can not be interpreted in geographical/ecological terms.

Latitude was also correlated with most prey items and trophic variables (Table 2).

FIGURE 2

Fish consumption showed a strong positive correlation with latitude, a pattern shared
20 by avian prey, while reptiles, crayfish and other aquatic invertebrates were more consumed at lower latitudes. The southward intensification of predation upon these

four prey categories occurred in parallel to an increase in diet diversity (Fig. 2B) and in the number of prey categories in otter diet, and a decrease in the number of fish families (Table 2). Predation upon mammals and amphibians did not show any geographical pattern. Diet diversity was negatively correlated with fish consumption and positively with the RFOs of amphibians, reptiles, crayfish and other aquatic invertebrates ($P < 0.001$ in all cases). The RFOs of mammals and birds was not related to diet diversity.

Two-sample comparisons between the otter diet in Mediterranean and temperate climatic areas produced similar results (Table 2). Mediterranean otters had more diverse diets and fed on a larger number of prey categories and a smaller number of fish families than those occupying temperate habitats. The frequency of occurrence of fish in otter diet was significantly higher in temperate than in Mediterranean locations, while the contrary happens with reptiles. On the average, crayfish and other invertebrates were consumed more often in Mediterranean habitats, while birds and mammals were more common prey in temperate ones, though these differences were not significant when corrected by Bonferroni's methods (Table 2). Differences between Mediterranean and temperate locations in the consumption of amphibians were smaller.

DISCUSSION

Though fish are the otter's main prey everywhere in Europe (Mason & Macdonald, 1986; Carrs, 1995; Kruuk, 1995), their consumption shows on average a marked decrease in southern localities. This is compensated with more intense

predation upon a pool of alternative prey including crayfish, reptiles, amphibians and aquatic invertebrates. Some of these alternative prey can constitute the bulk of otter diet in some southern places . In consequence, Mediterranean otters show higher diet diversity, predated upon a larger number of prey categories than those of temperate
5 habitats.

According to the latitudinal gradient in diversity, an increased trophic diversity of the otter in southern latitudes could be related to a higher abundance and availability of non-fish prey in these areas. In fact, at least reptiles and insects are specially abundant in Mediterranean Europe (Blondel & Aronson 1999), where the warm
10 weather conditions allow them to reach large sizes and to be active most of the year. Moreover, the recent spread of the introduced American crayfish (*Procambarus clarkii*, Girard) has changed the diet of otters and other predators in many Mediterranean habitats (Delibes & Adrián, 1987; Beja, 1996; Correia, 2001), crayfish becoming an important prey. However, a higher availability of alternative
15 prey makes possible, but not forces, an increase of trophic diversity. Besides this, it is usually necessary a reduction in the abundance or availability of the favourite prey (Stephens & Krebs, 1986). Erlinge (1968) stated that captive otters preferred to predate upon fish, and apparently fish are the otter's staple prey whenever abundant, even under Mediterranean conditions. In fact, otters have an almost exclusive
20 piscivorous diet in some Mediterranean localities where alternative prey are probably abundant (see Ruiz-Olmo *et al.*, 1989; Prigioni *et al.*, 1991). In the same way, otters in temperate areas feed on a high diversity of prey when availability of fish is reduced, like in Bialowieza, Poland (Brzeziński *et al.*, 1993; see Fig. 2). Kruuk

(1995) also related a increase of non-fish prey with periods of low fish abundance in Scotland rivers.

We suspect that the increased trophic diversity of Mediterranean otters is related also with a low or unpredictable availability of fish in freshwater Mediterranean ecosystems. The harsh environmental conditions during the strong summer drought are a key factor explaining the composition and dynamics of Mediterranean freshwater communities (Prenda & Gallardo, 1996). Most streams and small rivers become dry or broken in isolated pools during the summer, reducing the availability of fish (Pires *et al.*, 1999). Besides, the high intra and interannual variability in the precipitation and temperature regime characteristic of Mediterranean areas severely affects freshwater ecosystems, necessarily resulting in an unpredictable fish availability (Mooney, 1981; Prenda *et al.*, 2001). Thus, the widening of the otter's feeding niche in Mediterranean ecosystems would be favoured in an environment where fish populations are temporally scarce and patchily available, both in space and time (see Erlinge, 1986). Sulkava (1996) described a very diverse diet of otters in central Finland ($H' = 1.16$) (see Table 2), in a Taiga environment where the extremely cold winters produces great variations and unpredictability in fish availability (this work was not included in the analyses, since central Finland can not be considered a temperate area).

The review on otter diet by Jędrzejewska *et al.* (2001) did not find any relation between diversity or composition of otter diet and latitude, but concluded that otters behave as more generalist predators in streams and rivers than in lakes and sea shores, where fish are more frequent prey. This conclusion supports our hypothesis in an indirect way, as relates changes in otter's diet diversity with habitat features,

FIGURE 3

specially water availability and its effects on fish populations. We hypothesize that intra and interannual abundance and predictability of fish resources could range from a maximum in sea shores and lakes to a minimum in Mediterranean temporal streams, being intermediate in temperate water courses; this should be accompanied
5 by changes in the pattern of use of fish, its favourite prey, by the otter (Fig. 3).

Several reasons (criteria to select the dietary studies, covered range, oversight of including reptiles, etc) help to explain why Jędrzejewska *et al.* (2001) failed to describe the evident temperate-Mediterranean change in otter trophic niche breadth. However, the main cause probably was the low proportion of Mediterranean
10 localities included in their analysis: only 19 of 102 diet studies came from Mediterranean areas, and only 12 of them correspond to freshwater ecosystems. As the observed latitudinal pattern seems to be not gradual, but related to the different conditions in Mediterranean and temperate ecosystems, it could be concealed if the proportion of localities is very biased towards temperate places (see Fig. 2 to note
15 that in our study a latitudinal trend can be detected within Mediterranean localities, but is not apparent when only temperate ones are considered). Thus, Jędrzejewska *et al.* (2001) could detect the increase in otter trophic diversity from sea-shores and lakes to temperate rivers, but not the geographical pattern presented here, from temperate to Mediterranean freshwater habitats (Fig. 3).

20 The decrease in fish consumption and the corresponding enlargement of the food-niche of Mediterranean otters can be related with their small body size. Ruiz-Olmo *et al.* (1998) showed that otters from temperate areas were between 35% and 11% heavier than otters inhabiting the Iberian Peninsula. Some authors (King and Moody 1982, Clevenger, 1993) have related changes in mustelids body size with differences

in food abundance and availability. Temporally reduced fish abundance and unpredictable availability in freshwater Mediterranean environments could then favour small body sizes in otters, given that reduced energetic demands allow higher dietary flexibility (Gittleman & Purvis, 1998). Iriarte *et al.* (1990) also related the
5 reduced body size of American pumas in low latitudes with broader niche breaths.

The results of this paper show a clear dietary diversification of otters inhabiting Mediterranean freshwater ecosystems in relation with otters from temperate ones, resulting in a strong latitudinal gradient. Similar results have been published for other small and medium size predators in Mediterranean areas and explained also as a
10 response to the reduction in the diversity and abundance of their main prey (e.g. small mammals for the barn owl and the common genet; Herrera 1974, Virgós *et al.* 1999). This pattern can not be extended, however, to bigger Mediterranean predators, such as the Iberian Lynx (*Lynx pardinus*, Temminck), the golden eagle (*Aquila chrysaetos*, L.) (Delibes, 1975) and the badger (*Meles meles*, L.) (Martín *et al.*, 1995;
15 Goszczyński *et al.*, 2000), which in Mediterranean Iberia strongly predate upon rabbits (*Oryctogalus cuniculus*), an extremely abundant prey. Thus, a trend to an increased dietary niche breadth with reduced latitude can not be generalized.

ACKNOWLEDGMENTS

20 We thank Dr. B. Jędrzejewska, Dr. J. Jiménez Nieva, Dr. A. Rodríguez and Dr. J. Calzada for their helpful comments on early versions of the manuscript. The study was financially supported by GIASA-CSIC, through the project “Medidas compensatorias de la Autovía A-381 Jerez de la Frontera-Los Barrios”.

REFERENCES

- Acera, F. (1998). *Distribución y dieta de la nutria (Lutra lutra L., 1758) en el río Francia, Salamanca*. Tesis de Licenciatura, Universidad de Salamanca.
- Adrián, M. I. and Moreno, S. (1986). Notas sobre la alimentación de la nutria (*Lutra lutra*) en el embalse de Matavacas, Huelva. *Doñana, Acta Vertebrata*. **13**, 189-191.
- Adrián, M. I. and Delibes, M. (1987). Food habits of the otter (*Lutra lutra*) in two habitats of the Doñana National Park, SW Spain. *Journal of Zoology, London*. **212**, 399-406.
- Arcá, G. and Prigioni, C. (1987). Food of the Otter on the Fiora River (Central Italy). *Acta Theriologica*. **32**, 134-140.
- Arroyo, B. E. (1997). Diet of Montagu harrier *Circus pigargus* in Central Spain. Analysis of temporal and geographic variation. *Ibis*. **139**, 664-672.
- Beja, P. R. (1996). An analysis of otter *Lutra lutra* predation on introduced American crayfish *Procambarus clarkii* in Iberian streams. *J. Appl. Ecol.* **33**, 1156-1170
- Blondel, J. and Aronson, J. (1999). *Biology and wildlife of the Mediterranean Region*. Oxford University Press, Oxford.
- Brzeziński, M., Jędrzejewski, W. and Jędrzejewska, B. (1993). Diet of otters (*Lutra lutra*) inhabiting small rivers in the Bialowieza National Park, eastern Poland. *Journal of Zoology, London*. **230**, 495-501.
- Callejo, A. and Delibes, M. (1987). Dieta de la nutria *Lutra lutra* (Linnaeus, 1758) en la cuenca del alto Ebro, Norte de España. *Miscellanea Zoologica*. **11**, 353-362.

- Canas, C. (1999). *Predação por lontra, Lutra lutra L., em habitats estivais de uma ribeira intermitente Mediterrânica*. Relatório de estágio do curso de Licenciatura em Biologia Marinha e Pescas, Universidade do Algarve.
- Carss, D. N. (1995). Foraging behaviour and feeding ecology of the otter *Lutra lutra*:
5 a selective review. *Hystrix*. **7**, 179-194.
- Carss, D. N. and Parkinson, S. G. (1996). Errors associated with otter *Lutra lutra* faecal analysis. I. Assessing general diet from spraints. *Journal of Zoology, London*. **238**, 301-317.
- Carss, D. N., Kruuk, H. and Conroy, J. W. H. (1990). Predation on adult Atlantic
10 salmon, *Salmo salar* L., by otters, *Lutra lutra* (L.), within the River Dee system, Aberdeenshire, Scotland. *Journal of Fish Biology*. **37**, 935-944.
- Chanin, P. (1981). The diet of the otter and its relation with the feral mink in two areas of Southwest England. *Acta Theriologica*. **26**, 83-95.
- Clevenger, A. P. (1993). Pine marten (*Martes martes* Linné, 1758) comparative
15 feeding ecology in a island and mainland population of Spain. *Zeitschrift für Säugetierkunde*. **58**, 212-224.
- Correia, A. M. (2001). Seasonal and interspecific evaluation of predation by mammals and birds on the introduced red swamp crayfish *Procambarus clarkii* (Crustacea, Cambaridae) in a freshwater marsh (Portugal). *Journal of Zoology, London*. **255**, 533-541.
20
- Delibes, M. (1975). Some characteristic features of predation in the Iberian Mediterranean ecosystem. *XII Congresso da União Internacional dos Biologistas da Caça*, 31-36.

- Delibes, M. and Adrián, M. I. (1987). Effects of crayfish introduction on otter *Lutra lutra* food in the Doñana National Park, SW Spain. *Biol. Conserv.* **42**, 153-159.
- Emberger, L., Gaussen, H., Kassas, M. and de Philippis, A. (1963). *Carte bioclimatique de la zone méditerranéenne (Etude écologique de la zone méditerranéenne)*. UNESCO-FAO.
- 5 Erlinge, S. (1967). Food habits of the fish-otter *Lutra lutra* L. in South Swedish habitats. *Viltrevy.* **4**, 371-447.
- Erlinge, S. (1968). Food studies on captive otters *Lutra lutra* L. *Oikos.* **19**, 259-270.
- Erlinge, S. (1969). Food habits of the otter *Lutra lutra* L. and the mink *Mustela vison*
 10 Schreber in a trout water in southern Sweden. *Oikos.* **20**, 1-7.
- Erlinge, S. (1986). Specialists and generalists among mustelids. *Lutra.* **29**, 5-11.
- Futuyma, D. J. and Moreno, G. (1988). The evolution of ecological specialization. *Annual Review of Ecology and Systematics.* **19**, 207-233.
- Gasith, A and Resh, V. H. (1999). Streams in Mediterranean climate regions -
 15 Abiotic influences and biotic responses to predictable seasonal events. *Annual Review of Ecology and Systematics.* **30**, 51-81.
- Geidezis, L. (1998). What do otters (*Lutra lutra*) feed in a carp pond area in Saxony, Eastern Germany. *Boku-reports on wildlife research and game management.* **14**, 65-72.
- 20 Gittleman, J. L. & Purvis, A. (1998). Body-size and species-richness in carnivores and primates. *Proceedings of the Royal Society of London. Series B, Biological Sciences* **265**, 113-119.

- Gormally, M. J. and Fairley, J. S. (1982). Food of otters *Lutra lutra* in a freshwater lough and an adjacent brackish lough in the West of Ireland. *Journal of Zoology, London*. **197**, 313-321.
- Goszczyński, J., Jędrzejewska, B. and Jędrzejewski, W. (2000). Diet composition of badgers (*Meles meles*) in a pristine forest and rural habitats of Poland compared to other European populations. *Journal of Zoology, London*. **250**, 495-205.
- Hawkins, B. A. (2001). Ecology's oldest pattern?. *Trends in Ecology and Evolution*. **16**, 470.
- Henshilwood, D. A. (1981). *Status and diet of the otter (Lutra lutra) at Bosherton lakes, Dyfed*. Unpublished report. Cited in: Mason, C. F. and Macdonald, S. M. (1986). *Otters, ecology and conservation*. Cambridge: Cambridge University Press.
- Herrera, C. M. (1974). Trophic diversity of the Barn Owl *Tyto alba* in continental Western Europe. *Ornis Scandinavica* **5**, 181-191.
- Hulbert, S. H. (1984). Pseudoreplication and the design of ecological field experiments. *Ecological Monographs*. **54**, 187-211.
- Iriarte, A., Franklin, W. L., Johnson, W. E. & Redford, K. H. (1990). Biogeographic variation of food habits and body size of the American puma. *Oecologia*. **85**, 185-190.
- Jacobsen, L. and Hansen, H. M. (1996). Analysis of otter (*Lutra lutra*) spraints: Part 1: Comparison of methods to estimate prey proportions; Part 2: Estimation of the size of prey fish. *Journal of Zoology, London*. **238**, 167-180.

- Jędrzejewska, B., Sidorovich, V.E., Pikulik, M.M. and Jędrzejewski, W. (2001).
Feeding habits of the otter and the American mink in Białowieża Primeval Forest
(Poland) compared to other Eurasian populations. *Ecography*. **24**, 165-180.
- King, C. M. & Moody, J. E. (1982). The biology of the stoat (*Mustela erminea*) in
5 the National Parks of New Zealand III. Morphometric variation in relation to
growth, geographical distribution and colonisation. *New Zealand Journal of
Zoology*. **9**, 81-102.
- Knollseisen, M. and Kranz, A. (1998). Influence of different spraint sampling
methods on the results of otter diet studies. *Boku-reports on wildlife research and
10 game management*. **1**,: 37-45.
- Korpimäki, E. and Marti, C. D. (1995). Geographical trends in trophic characteristics
of mammal-eating and bird-eating raptors in Europe and North America. *Auk*.
112, 1004-1023.
- Kyne, M. J., Smal, C. M. And Fairley, J. S. (1989). The food of otters *Lutra lutra* in
15 the Irish Midlands and a comparison with that of the mink *Mustela vison* in the
same region. *Proceedings of the Royal Irish Academy*. **89**, 33-46.
- Kruuk, H. (1995). *Wild otters. Predation and populations*. Oxford University Press,
Oxford.
- Lanszki, J. and Körmendi, S. (1996). Otter diet in relation to fish availability in a fish
20 pond in Hungary. *Acta Theriologica*. **41**, 127-136.
- Libois, R. (1995). Régime et tactique alimentaires de la loutre (*Lutra lutra*) en
France: synthèse. *Cahiers d'Éthologie*. **15**, 251-274.

- Lodé, T. (1989). La loutre en Loire-Atlantique. Note préliminaire sur l'évolution récente des populations, l'alimentation et le comportement prédateur de *Lutra lutra*, L. 1758. *Bulletin de la Societe de Science Naturell de l'Ouest de la France*. **11**, 69-76.
- 5 López-Nieves, P. and Hernando, J. A. (1984). Food habits of the otter in central Sierra Morena (Córdoba, Spain). *Acta Theriologica*. **29**, 383-401.
- MacArthur, R. H. and Pianka, E. R. (1966). On optimal use of a patchy environment. *American Naturalist* **100**, 603-609.
- Martín, R., Rodríguez, A. and Delibes, M. (1995). Local feeding specialization by
10 badgers (*Meles meles*) in a Mediterranean environment. *Oecologia*. **101**, 45-50.
- Magalhães, M. F., Batalha, D. C. and Collares-Pereira, M. J. (2002). Gradients in stream fish assemblages across a Mediterranean landscape: contributions on environmental factors and spatial structure. *Freshwater Biology*. **47**, 1015-1031.
- Mason, C. F. and Macdonald, S. M. (1986). *Otters, ecology and conservation*.
15 Cambridge University Press, Cambridge.
- Mooney, H. A. (1981). Primary production in Mediterranean region. *Ecosystems of the World. Mediterranean type shrublands*. (ed. by di Castri, P., Goodwall, D. W. and Specht, R. L.). Elsevier.
- Morales, J. J. and Lizana, M. (1997). Autoecología y distribución de la nutria
20 euroasiática (*Lutra lutra* Linneo, 1758) en el Parque Natural del Lago de Sanabria y alrededores (Zamora). *Anuario Instituto de Estudios Zamoranos "Florián del Campo"*.: 339-395.

- Morales, J. J., Lizana, M., Gutiérrez, J. and Pedraza, E. (1998). *Distribución espacial y ecología trófica de la nutria euroasiática y el visón americano en el Parque Natural de Las Hoces del Río Duratón (Segovia)*. Obra Social y Cultural, Caja Segovia.
- 5 O'Neill, E. M., Day, K. and Paterson, J. P. H. (1998). Predation by otters at a salmon hatchery evidenced by diet studies. *Boku-reports on wildlife research and game management*. **14**, 46-64.
- Pires, A. M., Cowx, I. G. and Coelho, M. M. (1999). Seasonal changes in fish community structure in the middle reaches of the Guadiana basin, Portugal.
- 10 *Journal of Fish Biology*. **54**, 235-249.
- Prenda, J. and Gallardo, A. (1996). Self-purification, temporal variability and the macroinvertebrate community in small lowland Mediterranean streams receiving crude domestic sewage effluents. *Archiv für Hydrobiologie*, **136**, 159-170.
- Prenda, J., López-Nieves, P. and Bravo, R. (2001). Conservation of otter (*Lutra*
- 15 *lutra*) in a Mediterranean area: the importance of habitat quality and temporal variation in water availability. *Aquatic Conservation: Marine and Freshwater Ecosystems*. **11**, 343-355.
- Prigioni, C., Pandolfi, M., Grimod, I., Fumagalli, R., Santolini, R., Arcá, G., Montemurro, F., Bonacoscia, M. and Racana, A. (1991). The otter in five Italian
- 20 rivers.- First report. *Proceedings V International Otter Colloquium, Hankensbüttel 1989, Habitat*. **6**, 143-145.
- Rosenzweig, M. L. (1996). *Species diversity in space and time*. Cambridge University Press, Cambridge.

- Ruiz-Olmo, J. (1995). The reptiles in the diet of the otter (*Lutra lutra* L., Carnivora, Mammalia) in Europe. *Scientia Herpetologica*. **1**, 259-264.
- Ruiz-Olmo, J., Delibes. M. and Zapata, S.C. (1998). External morphometry, demography and mortality of the otter *Lutra lutra* (Linneo, 1758) in the Iberian Peninsula. *Galemys*. **10**, 239-251.
- Ruiz-Olmo, J., Jordán, G. and Gosálbez, J. (1989). Alimentación de la nutria (*Lutra lutra* L., 1758) en el Nordeste de la Península Ibérica. *Doñana, Acta Vertebrata*. **16**, 227-237.
- Schoener, T. W. (1971). Theory of feeding strategies. *Annual Review of Ecology and Systematics*. **2**, 369-404.
- Smith, R. L. and Smith, T. M. (2000). *Elements of ecology*, 4th ed. Addison Wesley Longman, Inc.
- Sokal, R. R. and Rohlf, F. J. (1995). *Biometry*. W.H. Freeman and Company, New York.
- Stephens, D. W. and Krebs, J. R. (1986). *Foraging theory*. Monographs in behaviour and ecology. Princeton University Press, Princeton.
- Sulkava, R. (1996). Diet of otters *Lutra lutra* in central Finland. *Acta Theriologica*. **41**, 395-408.
- Taastrøm, H. M. and Jacobsen, L. (1999). The diet of otters (*Lutra lutra* L.) in Danish freshwater habitats: comparisons of prey fish populations. *Journal of Zoology, London*. **248**, 1-13.
- Virgós, E., Llorente, M. and Cortés, Y. (1999). Geographical variation in genet (*Genetta genetta* L.) diet: a literature review. *Mammal Review*. **29**, 119-128.

- Webb, J. B. (1975). Food of the otter (*Lutra lutra*) on the Somerset levels. *Journal of Zoology, London*. **177**, 486-491.
- Weber, J. M. (1990). Seasonal exploitation of amphibians by otters (*Lutra lutra*) in north-east Scotland. *Journal of Zoology, London*. **220**, 641-651.
- 5 Wise, M. H., Linn, I. J. and Kennedy, C. R. (1981). A comparison of the feeding biology of mink *Mustela vison* and otter *Lutra lutra*. *Journal of Zoology, London*. **195**, 181-213.
- Wisniowska, L. (1996). Diet of the otter (*Lutra lutra*) in fish ponds in Southern Poland. *Journal of Wildlife Research*. **1**, 272-277.
- 10 Zar, J. H. (1984). *Biostatistical analysis*, 2nd edn. Prentice Hall.

Table 1. Mean Relative Frequency of Occurrence (RFO) for several otter (*Lutra lutra* L.) prey items in Europe and Pearson's correlation (r) between them and the principal component 1. Significance levels: ***, P< 0.001; **, P< 0.01; *, P< 0.05., NS, non significant.

5

Prey items	Mean \pm SD	r
		Factor 1
Fish	72.8 \pm 18.5	0.91***
Amphibians	8.8 \pm 7.6	-0.68***
Reptiles	1.5 \pm 2.9	-0.61***
Birds	1.9 \pm 2.1	0.31 (NS)
Mammals	0.9 \pm 1.3	0.43**
Crayfish	6.8 \pm 12.9	-0.49**
Aq. invertebrates	7.0 \pm 7.7	-0.79***
<i>Eigenvalue</i>		2.80
<i>% Explained Variance</i>		40.0

Table 2. Pearson (*r*) significant correlations ($P \leq 0.05$) between latitude and RFOs of general prey items in otter diet, diet diversity (H'), number of prey items in otter diet (NPC) and number of fish families in otter diet (NFF). Significance levels for *r* values as for table 1. Comparisons of the different variables under different climatic conditions using t-test are also shown. Significance levels for marked (*) *t* values, $P < 0.005$ (Bonferroni's correction; $P = 0.05/10$ tests).


	Latitude	t-test results		
		Mean \pm SD		
		Temperate (n= 21)	Mediterranean (n= 16)	<i>t</i>
<i>r</i>				
Fish	0.51***	80.6 \pm 15.5	62.6 \pm 17.5	3.36*
Amphibians	-0.16 (NS)	8.2 \pm 9.0	9.6 \pm 5.3	1.01
Reptiles	-0.71***	0.1 \pm 0.2	3.3 \pm 3.7	6.82*
Birds	0.36*	2.6 \pm 2.3	1.0 \pm 1.2	2.77
Mammals	0.16 (NS)	1.1 \pm 1.3	0.7 \pm 1.3	1.61
Crayfish	-0.38*	2.9 \pm 6.8	11.8 \pm 17.0	2.41
Aq. invertebrates	-0.51***	4.3 \pm 5.9	10.5 \pm 8.7	2.64
H'	-0.57***	0.62 \pm 0.35	0.98 \pm 0.28	3.62*
NPC	-0.55***	5.1 \pm 1.15	6.2 \pm 0.58	3.51*
NFF	0.56***	5.3 \pm 1.38	3.7 \pm 0.79	4.18*


FIGURE CAPTIONS

- Figure 1.** Distribution of the 37 reviewed studies on the diet of otters (*Lutra lutra*, L.). The dotted line separates temperate (filled circles) from Mediterranean locations (empty circles). The numbers are assigned to original works as follows. 1: Canas (1999); 2: Beja (1996); 3: Adrián and Moreno (1986); 4, 5: Adrián and Delibes (1987); 6: López-Nieves and Hernando (1984); 7: Morales and Lizana (1997); 8: Acera (1998); 9: Morales et al. (1998); 10: Callejo and Delibes (1987); 11, 12: Ruiz-Olmo et al. (1989); 13: Arcá and Prigioni (1987); 14, 15, 16: Prigioni et al. (1991); 17: Lóde (1989); 18: Libois (1995); 19, 20: Chanin (1981); 21: Wise et al. (1981); 22: Webb (1975); 23: Henshilwood (1981); 24: Gormally and Fairley (1982); 25: Kyne et al. (1989); 26: O'Neill et al. (1998); 27, 29: Weber (1990); 28: Carss et al. (1990); 30: Taastrøm and Jacobsen (1999); 31: Erlinge (1967); 32: Erlinge (1969); 33: Geidezis (1998); 34: Lanszki and Körmendi (1996); 35: Knollseisen and Kranz (1998); 36: Wisniowska (1996); 37: Brzeziński et al. (1993)..
- Figure 2.** Relation between latitude and: A) principal component 1 (PC 1) scores ($r=0.63$; $P<0.001$) and B) diet diversity ($r=-0.57$; $P<0.001$) for 37 otter (*Lutra lutra*) diet studies. Filled circles: temperate locations; empty circles: Mediterranean locations. Encircled point (Brzeziński *et al.*, 1993; point 37 Fig. 1) is commented in Discussion.
- Figure 3.** Schematic diagram representing the suggested variation in otter diet diversity and fish communities characteristics in Europe in relation to water availability and stability. An amplification in otter diet diversity is observed as fish become scarce or unpredictably available. Numbers show the habitats in which the


review by Jędrzejewska *et al.* (2001) (1) and this study (2) were centred.

Clavero *et al.* Changes in otter diet diversity in Europe. FIGURE 1



Clavero *et al.* Changes in otter diet diversity in Europe. FIGURE 2

Clavero *et al.* Changes in otter diet diversity in Europe. FIGURE 3



BIOSKETCHES

Miguel Clavero had a grant from the Doñana Biological Station, from the Spanish Council for Research (CSIC), and at present is an associate professor in the Department of Environmental Biology and Public Health in Huelva University. He is
5 preparing his PhD thesis on the ecology and conservation of the otter in freshwater and coastal habitats.

José Prenda is an assistant professor in the Department of Environmental Biology and Public Health in Huelva University. His research works have been centred in the ecology and conservation problems of Mediterranean river ecosystems,
10 including macroinvertebrates, fishes and the otter.

Miguel Delibes is a research professor in the Department of Applied Biology of the Doñana Biological Station, from the Spanish Council for Research (CSIC). He is a world known conservation biologist and an expert in the ecology of Mediterranean carnivores.