

Communications: Evidence for the role of fluctuations in the thermodynamics of nanoscale drops and the implications in computations of the surface tension

José G. Sampayo, Alexandr Malijevský, Erich A. Müller, Enrique de Miguel, and George Jackson

Citation: *The Journal of Chemical Physics* **132**, 141101 (2010); doi: 10.1063/1.3376612

View online: <http://dx.doi.org/10.1063/1.3376612>

View Table of Contents: <http://scitation.aip.org/content/aip/journal/jcp/132/14?ver=pdfcov>

Published by the [AIP Publishing](#)

Articles you may be interested in

[Surface tension of spherical drops from surface of tension](#)

J. Chem. Phys. **140**, 034110 (2014); 10.1063/1.4862149

[A fingerprint of surface-tension anisotropy in the free-energy cost of nucleation](#)

J. Chem. Phys. **138**, 064508 (2013); 10.1063/1.4790635

[Statics of polymer droplets on deformable surfaces](#)

J. Chem. Phys. **135**, 214703 (2011); 10.1063/1.3663381

[On the interfacial thermodynamics of nanoscale droplets and bubbles](#)

J. Chem. Phys. **135**, 024701 (2011); 10.1063/1.3609274

[Discussion on a mechanical equilibrium condition of a sessile drop on a smooth solid surface](#)

J. Chem. Phys. **130**, 144106 (2009); 10.1063/1.3103269



Communications: Evidence for the role of fluctuations in the thermodynamics of nanoscale drops and the implications in computations of the surface tension

José G. Sampayo,¹ Alexandr Malijevský,^{1,2} Erich A. Müller,¹ Enrique de Miguel,³ and George Jackson^{1,a)}

¹Department of Chemical Engineering, Imperial College London, South Kensington Campus, London SW7 2AZ, United Kingdom

²E. Hála Laboratory of Thermodynamics, Institute of Chemical Process Fundamentals of ASCR, 16502 Prague 6, Czech Republic

³Departamento de Física Aplicada, Facultad de Ciencias Experimentales, Universidad de Huelva, 21071 Huelva, Spain

(Received 27 January 2010; accepted 10 March 2010; published online 9 April 2010)

Test-area deformations are used to analyze vapor-liquid interfaces of Lennard-Jones particles by molecular dynamics simulation. For planar vapor-liquid interfaces the change in free energy is captured by the average of the corresponding change in energy, the leading-order contribution. This is consistent with the commonly used mechanical (pressure-tensor) route for the surface tension. By contrast for liquid drops, one finds a large second-order contribution associated with fluctuations in energy. Both the first- and second-order terms make comparable contributions, invalidating the mechanical relation for the surface tension of small drops. The latter is seen to increase above the planar value for drop radii of ~ 8 particle diameters, followed by an apparent weak maximum and slow decay to the planar limit, consistent with a small negative Tolman length. © 2010 American Institute of Physics. [doi:10.1063/1.3376612]

It is striking that though almost a century has passed since Gibbs formulated his thermodynamic theory of curved interfaces, there is still widespread controversy about the dependence of the surface tension on the curvature (size of a drop) and the validity of the mechanical route to the surface tension.¹⁻³ The formal approach of Gibbs is intimately connected with the relations of Laplace, $\Delta p = 2\gamma_s/R_s$, and Tolman, $\gamma(R)/\gamma_\infty = 1 - 2\delta_\infty/R + \dots$, for drops of radius R . Here, $\Delta p = p_l - p_g$ is the pressure difference inside (l) and outside (g) the drop, $\gamma_s = \gamma(R_s)$ is the interfacial tension associated with the surface of tension R_s , γ_∞ is the value for the planar gas-liquid surface, and the Tolman⁴ length δ_∞ is defined relative to the radius of the equimolar surface R_e as $\delta_\infty = \lim_{R_s \rightarrow \infty} (R_e - R_s)$.

There are three basic routes to the definition of the tension:¹ Thermodynamic (Gibbs and Tolman), mechanical (Laplace and Young), and statistical mechanical (density functional and related theories). The thermodynamic and mechanical routes are macroscopic theories, so there has been much debate about their applicability to small systems such as nanoscale liquid drops or bubbles. One key question is whether the mechanical relations based on the pressure virial (formulated in terms of the appropriate tensorial components) that make use of the concept of the bulk pressure of the coexisting states are appropriate at these length scales for curved surfaces.

While the Laplace equation essentially defines the ratio γ_s/R_s , the first-order form of Tolman's theory is appropriate

only for sufficiently large drops. One can view δ_∞ as the leading-order correction to the tension of a planar surface. Despite its fundamental role in studies of interfacial properties of curved surfaces and theories of nucleation, there is still much controversy as to even the *sign* of δ_∞ . Microscopic statistical mechanical approaches including square gradient theories (SGTs),^{5,6} curvature expansions of the planar interface,^{7,8} and density functional theories (DFTs), including local⁹⁻¹¹ and nonlocal¹²⁻¹⁴ treatments, have led to conflicting views on the magnitude and sign of δ_∞ , as well as the curvature dependence of the surface tension. The widely accepted view from this body of work is that $\delta_\infty \leq 0$ and that there is a small maximum in $\gamma_s(R)$ as the drop radius is decreased, then followed by a sharp decrease. This is supported by studies on the penetrable sphere model¹⁵ (which can be solved exactly at the mean-field level at zero temperature) where one finds a negative Tolman length ($\delta_\infty = -\sigma/2$), with σ the molecular diameter.

By contrast, the vast majority of computer simulation studies suggests that $\delta_\infty > 0$. In most simulations of liquid drops, the mechanical route to the interfacial tension is employed, usually involving an integration of the gradient of the normal component of the pressure tensor from the center of the drop to the bulk vapor phase.¹⁶⁻¹⁹ In this case one predicts a monotonous decrease in the surface tension with increasing curvature (decreasing drop radius) from the planar limit (infinite radius); this would correspond to $\delta_\infty > 0$ throughout. As was pointed out early on by Schofield and Henderson,² there are fundamental problems in employing local pressure tensors and the associated definition of the internal pressure for microscopic (high curvature) drops.

^{a)}Author to whom correspondence should be addressed. Electronic mail: g.jackson@imperial.ac.uk.

This leads to a mismatch in the free energy of the formation of a drop determined via the mechanical and thermodynamic routes as observed in simulation.²⁰ Macroscopic thermodynamic routes based on a combination of the Laplace and Tolman relations have been employed¹⁶ but also suffer from the ill-definition of the internal pressure and density of the liquid. One can estimate the interfacial tension from the free energy change accompanying a volume deformation of spherical surfaces using a virial-like expression;²¹ these results for the surface tension are in disagreement with those obtained from the direct mechanical route. Recent grand canonical simulations²² and a thermodynamic analysis of large drops based on the Laplace–Tolman relations²³ both now appear to suggest a small negative δ_∞ , which is consistent with the findings of DFT.

The aim of this paper is to use a new method for the calculation of the surface tension of small liquid drops in molecular simulation, highlighting the role played by the fluctuations in the energy of deformation. The method relies on the thermodynamic definition of the surface tension and is thus free from the inconsistencies associated with the application of the mechanical route. A variant of the test-area (TA) method²⁴ is used where small virtual perturbations are made in the box dimensions of systems with interfaces to obtain the change in free energy associated with the corresponding change in surface area. For a fluid drop of radius R , the change in the Helmholtz free energy F is expressed thermodynamically as¹

$$dF = -SdT - p_g dV_g + p_l dV_l + \mu dN + \gamma dA + CdR, \quad (1)$$

where S is the entropy, $V_{g,l}$ are the vapor and liquid volumes, T is the temperature, μ is the chemical potential, N is the number of particles, A is the interface area, and C is the conjugate variable for R . The surface tension of a drop is given by

$$\left(\frac{\partial F}{\partial A}\right)_{NVT} = \gamma_s, \quad (2)$$

where the minimal interfacial tension γ_s defines R_s and corresponds to taking $C=0$. The change in free energy ΔF due to a virtual change in area ΔA can be expressed as the average of the Boltzmann factor of the corresponding change in configurational energy ΔU ,²⁴

$$\Delta F = -kT \ln \left\langle \exp \left(-\frac{\Delta U}{kT} \right) \right\rangle \quad (3)$$

$$= \langle \Delta U \rangle - \frac{1}{2kT} \{ \langle \Delta U^2 \rangle - \langle \Delta U \rangle^2 \} + \frac{1}{6(kT)^2} \{ \langle \Delta U^3 \rangle - 3\langle \Delta U^2 \rangle \langle \Delta U \rangle + 2\langle \Delta U \rangle^3 \}. \quad (4)$$

The averages are over configurations of the unperturbed reference system. In Eq. (4) ΔF is expressed as a perturbation series to $\mathcal{O}(\langle \Delta U^3 \rangle)$, where the first-order average of the change in energy is $\Delta F_1 = \langle \Delta U \rangle$, the second-order energy fluctuation term is $\Delta F_2 = -\{ \langle \Delta U^2 \rangle - \langle \Delta U \rangle^2 \} / (2kT)$, and the third-order contribution is denoted by ΔF_3 . The full Boltz-




FIG. 1. TA deformations of a planar liquid-vapor interface of the LJ-TS fluid. MD simulations of $N=749$ particles in a periodic box of dimensions $L_x=L_y=7.885\sigma$ and $L_z=6L_x$ at $T^*=kT/\epsilon=0.8$ over 3×10^6 timesteps. The deformations correspond to changes in the box dimensions (particle coordinates) of $L'_x=L_x\sqrt{1+\xi}$, $L'_y=L_y\sqrt{1+\xi}$, and $L'_z=L_z/(1+\xi)$. (a) The contributions $\Delta F_1/\Delta A$ and $\Delta F_2/\Delta A$ to the change in free energy per unit area (in units of ϵ/σ^2) ($\Delta A^* < 0$, +; $\Delta A^* > 0$, ×; and average, ●). The interfacial tension $\gamma^* = \gamma\sigma^2/\epsilon$ is obtained by extrapolation to $\Delta A^* = \Delta A/\sigma^2 = 0$. (b) The distribution $\mathcal{P}(\Delta U)$ of the change in energy (relative to its average in units of ϵ) scaled at the maximum peak height for different relative deformations ΔA^* . The width (standard deviation, $\sigma_{\Delta U}$) is depicted in the inset.

mann form, Eq. (3), is employed in, e.g., the test-particle approach for the chemical potential,²⁵ or the volume perturbation method for the pressure²⁶ and the pressure tensor.²⁷

The tension is obtained as the change in free energy per unit area for infinitesimal perturbations to $\mathcal{O}(\langle \Delta U^3 \rangle)$,

$$\gamma = \lim_{\Delta A \rightarrow 0} \frac{\Delta F}{\Delta A} = \lim_{\Delta A \rightarrow 0} \left\{ \frac{\Delta F_1}{\Delta A} + \frac{\Delta F_2}{\Delta A} + \frac{\Delta F_3}{\Delta A} \right\}. \quad (5)$$

The leading term, $\Delta F_1 = \langle \Delta U \rangle$, corresponds to the mechanical work involved in changing the area of the interface, which can be directly associated with the so-called virial expression for the tension²⁸ (expressed in terms of averages of the appropriate components α of the virial, $\langle x_\alpha (dU/dx_\alpha) \rangle$, at the Hookean linear-response level). The corresponding entropic contribution due to the deformation is²⁸ $T\Delta S = \{ \langle U \rangle \langle \Delta U \rangle - \langle U \Delta U \rangle \} / (kT)$.

In the case of a planar interface, it is well known that the interfacial tension can be obtained formally from the virial expression,^{1,28} i.e., entirely from the leading-order contribution of Eq. (5). This is exemplified for a planar vapor-liquid interface of Lennard-Jones (LJ) particles (of diameter σ and well depth ϵ , truncated and shifted TS at $r_c=2.5\sigma$) as shown in Fig. 1. A planar interface is first stabilized during an NVT molecular dynamics (MD) simulation of the inhomogeneous system with a liquid slab in the center of a box




FIG. 2. TA ellipsoidal deformations of a spherical drop of LJ-TS liquid of radius $\langle R_e \rangle = 5.55\sigma$ in coexistence with its vapor. MD simulations of $N = 749$ particles in a periodic box of dimensions $L_x = L_y = L_z = 20\sigma$ at $T^* = 0.8$ over 1.5×10^9 timesteps. The deformations correspond to changes in the box dimensions (particle coordinates) of $L'_i = L_i \sqrt{1 + \xi}$, $L'_j = L_j \sqrt{1 + \xi}$, and $L'_k = L_k / (1 + \xi)$ (where i, j , and k denote any of the Cartesian axes). (a) The contributions $\Delta F_1/\Delta A$, $\Delta F_2/\Delta A$, and $\Delta F_3/\Delta A$ to the change in free energy per unit area (in units of ϵ/σ^2) (prolate, +; oblate, \times ; and average, \bullet). The tension $\gamma^* = \gamma\sigma^2/\epsilon$ is obtained by extrapolation to $\Delta A^* = \Delta A/\sigma^2 = 0$. (b) The distribution $\mathcal{P}(\Delta U)$ of the change in energy scaled at the maximum peak height for different relative deformations ΔA^* ; the width $\sigma_{\Delta U}$ is depicted in the inset. The Gaussians for the planar interface are shown dotted (note the very small scale in comparison).

separated by two vapor regions. The change in configurational energy due to small test changes in the dimensions of the box such that the interfacial area is increased or decreased at fixed overall volume is then computed to estimate the various contributions in Eq. (5); the limit of infinitesimal deformations is obtained by extrapolation to $\Delta A \rightarrow 0$. From Fig. 1(a) it is clear that only the leading *mechanical* term $\Delta F_1/\Delta A$ contributes to the interfacial tension of a planar interface, confirming the validity of the pressure-tensor route in this case. The fluctuation term $\Delta F_2/\Delta A$ is very small by comparison and does not contribute to the tension in an appreciable way; this is also true for the third-order term. In Fig. 1(b) we plot the distribution $\mathcal{P}(\Delta U)$ of the change in configurational energy (relative to $\langle \Delta U \rangle$) for different area perturbations; the distribution is well represented by a Gaussian, the width of which ($\sigma_{\Delta U}$) decreases to zero with $\Delta A \rightarrow 0$, consistent with a very small $\Delta F_2/\Delta A \sim 1 \times 10^{-6} \epsilon/\sigma^2$.

The overall physical picture is fundamentally different for a nanosized spherical drop of liquid in contact with its vapor. Once the drop has been stabilized, its size can be characterized from the density profile $\rho(r)$ as a function of the distance r from its center by calculating the Gibbs dividing surface $R_e^3 = (\rho_v - \rho_l)^{-1} \int dr r^3 d\rho(r)/dr$, corresponding to an area of $A = 4\pi R_e^2$. Virtual perturbations from the equilibrium spherical drop geometry are made with test changes in the




FIG. 3. The surface tension of spherical drops of LJ-TS fluids with average radii $\langle R_e \rangle = 5.55, 6.12, 8.04, 11.7$, and 14.6 at $T^* = 0.8$ from TA ellipsoidal deformations (\bullet), compared with the values from the mechanical route (Ref. 19) (\square), and the data of Thomson *et al.* (Ref. 16) (\triangle), El Bardouni *et al.* (Ref. 21) (\diamond), and Schrader *et al.* (Ref. 22) (continuous curve); the planar limit is shown dotted. The predictions of FMT (Ref. 14) are depicted in the inset (dashed curve).

dimensions of the simulation cube: Two of the Cartesian axes are decreased (or increased) in length, and the third is increased (or decreased) such that the overall volume remains constant. The perturbed states correspond to ellipsoidal drops of prolate (or oblate) shape, which always have larger surface areas than the original drop, $\Delta A > 0$. This essentially corresponds to the longest P_2 (Legendre polynomial) capillary-wave oscillations possible for the drop;³ the capillary-wave surface tension is equivalent to the thermodynamic one at least to leading order in curvature $\mathcal{O}(1/R)$. Averages are then accumulated over very long runs of $\sim 1.5 \times 10^9$ timesteps, corresponding to microsecond runs for typical molecular parameters. The term $\Delta F_1/\Delta A$ is more than two orders of magnitude larger in the case of the drop than for the planar interface system of comparable size [cf. Figs 1(a) and 2(a)]. The most significant difference is the large contribution from the second-order energy “fluctuation” term $\Delta F_2/\Delta A$ for the drop, which was negligible for the planar interface; this term is now comparable in magnitude to, but of opposite sign than, the first-order term. The third-order terms remain essentially negligible. As a result, both the first- and second-order terms contribute to the surface tension of the drop. A thermodynamic characteristic of the drop is thus the nonvanishing (and large) fluctuation term, which is clearly an indication of an additional entropic contribution. This can be seen in the distribution of the change in configurational energy for different TA perturbations [Fig. 2(b)]. The data are again well described as Gaussians, but though the width now appears not to vanish in the limit $\Delta A \rightarrow 0$ its variance of course does, i.e., $\lim_{\Delta A \rightarrow 0} \sigma_{\Delta U}^2 = 0$ and where in this case $\lim_{\Delta A \rightarrow 0} \Delta F_2/\Delta A \neq 0$. The fact that $\lim_{\Delta A \rightarrow 0} \Delta F_3 \sim 0$ for both the planar and curved systems suggests symmetrical Gaussians.

The dependence of the surface tension computed from ellipsoidal deformations as a function of the drop size (for systems with $N = 749$ to $11\,334$) is depicted in Fig. 3. Here the tension is computed for R_e rather than R_s though $\gamma_e = \gamma_s$ to $\mathcal{O}(1/R^2)$. The behavior obtained with our thermodynamic TA approach does not support the findings obtained from a standard pressure-tensor route (e.g., the recent MD data of

Vrabec *et al.*¹⁹). This is in line with the concerns of Schofield and Henderson,² and others^{8,20,29} about the inadequacy of the mechanical route for very small systems. For drops larger than $R_e \sim 8$ we observe values of the surface tension which appear to be slightly larger than the planar limit, $\gamma(R) > \gamma_\infty$; because the tension has to converge to γ_∞ when $R \rightarrow \infty$, this suggests a nonmonotonic behavior of the tension with increasing curvature and a corresponding weak maximum. Our values are consistent with the data point reported by El Bardouni *et al.*²¹ estimated from the surface free energy change and with the small maximum observed by Schrader *et al.*²² using a Landau free energy approach in the canonical ensemble (though the authors do not comment explicitly on this point). The calculations of the tension of curved interfaces from curvature corrections, SGT, and DFT (which have been brought into question because of their failure to reproduce existing simulation data) are now supported by our data. In the inset of Fig. 3 we compare the TA data for the surface tension of drops with those from a nonlocal DFT using fundamental measure theory (FMT);¹⁴ a maximum is predicted with FMT at $R_e \sim 10$.

Three main conclusions can be gleaned from our study. First, there is clearly a large fluctuation contribution to the interfacial tension of nanoscale spherical drops (and most likely other curved surfaces) in addition to the underlying first-order (mechanical) contribution, which fully describes the planar interface. Such contributions from fluctuations in the energy are not found in the planar limit to any significant degree (at least for the conditions examined here away from the critical region). Second, our results do not therefore support the validity of a mechanical (pressure-tensor) route to the interfacial tension for surfaces of high curvature such as small drops. This is in line with the warning of Blokhuis and Bedeaux⁸ that the use of a mechanical approach in this context “is still a matter of concern” and that it is “advisable not to use the pressure-tensor whenever this can be avoided.” Our data are not consistent with the monotonic dependence of the surface tension with curvature obtained from a mechanical treatment. As well as contributions in $\langle x_\alpha(dU/dx_\alpha) \rangle$, the correct “virial” expression for the surface tension would have to contain terms in averages of the type $\langle x_\alpha(dU/dx_\alpha) \rangle \times \langle x_\beta(dU/dx_\beta) \rangle$ and $\langle x_\alpha x_\beta(dU/dx_\alpha)(dU/dx_\beta) \rangle$, which would involve up to four-body correlations for pairwise additive potentials. This suggests that there are additional contributions to the change in the entropy due to the deformation of small drops involving quadratic terms in ΔU : $\langle \Delta U^2 \rangle$, $\langle \Delta U \rangle^2$, $\langle U \Delta U^2 \rangle$, $\langle \Delta U \rangle \langle U \Delta U \rangle$, $\langle U \rangle \langle \Delta U^2 \rangle$, and $\langle U \rangle \langle \Delta U \rangle^2$. As a final point, the rise in the surface tension above that of the planar limit after a certain drop size would be consistent with a negative Tolman length. Our data for the larger drops suggest a value of $\delta_\infty/\sigma \sim -0.2 \pm 0.3$. Though the statistical uncer-

tainty is large, our finding supports the exact mean-field predictions for the penetrable sphere model¹⁵ and is consistent with the latest accurate value of -0.10 ± 0.02 determined from the Laplace relation for a large $N=100\,000$ particle system.²³

We are very appreciative to Jim Henderson for useful discussions. J.G.S. acknowledges financial support from the CONACYT of Mexico for a Ph.D studentship, A.M. from the GACR (Grant No. IAA200760905) and GAAS (Grant No. IAA400720710) of the Czech Republic, and E.d.M. from the DGI (Grant No. FIS2007-66079-C02-02) and the Junta de Andalucía (Grant No. P07-FQM02884) of Spain. We also acknowledge funding from the EPSRC of the United Kingdom (Grant No. EP/E016340).

¹J. S. Rowlinson and B. Widom, *Molecular Theory of Capillarity* (Oxford University Press, Oxford, 1982).

²P. Schofield and J. R. Henderson, Proc. R. Soc. London **A379**, 231 (1982).

³J. R. Henderson, in *Fluid Interfacial Phenomena*, edited by C. A. Croxton (Wiley, New York, 1986).

⁴R. C. Tolman, *J. Chem. Phys.* **17**, 333 (1949).

⁵A. H. Falls, L. E. Scriven, and H. T. Davis, *J. Chem. Phys.* **75**, 3986 (1981).

⁶R. Guermeur, F. Biquard, and C. Jacolin, *J. Chem. Phys.* **82**, 2040 (1985).

⁷E. M. Blokhuis and D. Bedeaux, *Physica A* **184**, 42 (1992).

⁸E. M. Blokhuis and D. Bedeaux, *J. Chem. Phys.* **97**, 3576 (1992).

⁹D. J. Lee, M. M. Telo da Gama, and K. E. Gubbins, *J. Chem. Phys.* **85**, 490 (1986).

¹⁰D. W. Oxtoby and R. Evans, *J. Chem. Phys.* **89**, 7521 (1988).

¹¹K. Koga, X. C. Zeng, and A. K. Shchekin, *J. Chem. Phys.* **109**, 4063 (1998).

¹²T. V. Bykov and X. C. Zeng, *J. Chem. Phys.* **117**, 1851 (2002).

¹³Z. Li and J. Wu, *Ind. Eng. Chem. Res.* **47**, 4988 (2008).

¹⁴A. Malijevský and G. Jackson (unpublished).

¹⁵S. J. Hemingway, J. R. Henderson, and J. S. Rowlinson, *Faraday Symp. Chem. Soc.* **16**, 33 (1981).

¹⁶S. M. Thompson, K. E. Gubbins, J. P. R. B. Walton, R. A. R. Chantry, and J. S. Rowlinson, *J. Chem. Phys.* **81**, 530 (1984).

¹⁷M. J. P. Nijmeijer, C. Bruin, A. B. van Woerkom, A. F. Bakker, and J. M. J. van Leeuwen, *J. Chem. Phys.* **96**, 565 (1992).

¹⁸Y. A. Lei, T. Bykov, S. Yoo, and X. C. Zeng, *J. Am. Chem. Soc.* **127**, 15346 (2005).

¹⁹J. Vrabec, G. K. Keddia, G. Fuchs, and H. Hasse, *Mol. Phys.* **104**, 1509 (2006).

²⁰P. R. ten Wolde and D. Frenkel, *J. Chem. Phys.* **109**, 9901 (1998).

²¹H. El Bardouni, M. Mareschal, R. Lovett, and M. Baus, *J. Chem. Phys.* **113**, 9804 (2000).

²²M. Schrader, P. Virnau, and K. Binder, *Phys. Rev. E* **79**, 061104 (2009).

²³A. E. van Giessen and E. M. Blokhuis, *J. Chem. Phys.* **131**, 164705 (2009).

²⁴G. J. Gloor, G. Jackson, F. J. Blas, and E. de Miguel, *J. Chem. Phys.* **123**, 134703 (2005).

²⁵B. Widom, *J. Chem. Phys.* **39**, 2808 (1963).

²⁶R. Eppenga and D. Frenkel, *Mol. Phys.* **52**, 1303 (1984).

²⁷E. de Miguel and G. Jackson, *J. Chem. Phys.* **125**, 164109 (2006).

²⁸J. Lekner and J. R. Henderson, *Mol. Phys.* **34**, 333 (1977).

²⁹H. Reiss and D. Reguera, *J. Phys. Chem. B* **108**, 6555 (2004).